



RESEARCH ARTICLE

OPEN ACCESS

Modelling of the leaf area for various pear cultivars using neuro computing approaches

Ahmet Öztürk (Ozturk, A)¹, Bilal Cemek (Cemek, B)², Hüsnü Demirsoy (Demirsoy, H)¹ and Erdem Küçüktopcu (Kucuktopcu, E)²

¹Ondokuz Mayıs University, Faculty of Agriculture, Dept. Horticulture. 55139 Samsun, Turkey. ²Ondokuz Mayıs University, Faculty of Agriculture, Dept. Agricultural Structures and Irrigation. 55139 Samsun, Turkey.

Abstract

Aim of study: Leaf area (LA) is an important variable for many stages of plant growth and development such as light interception, water and nutrient use, photosynthetic efficiency, respiration, and yield potential. This study aimed to determine the easiest, most accurate and most reliable LA estimation model for the pear using linear measurements of leaf geometry and comparing their performance with artificial neural networks (ANN).

Area of study: Samsun, Turkey.

Material and methods: Different numbers of leaves were collected from 12 pear cultivars to measure leaf length (L), and width (W) as well as LA. The multiple linear regression (MLR) was used to predict the LA by using L and W. Different ANN models comprising different number of neuron were trained and used to predict LA.

Main results: The general linear regression LA estimation model was found to be $LA = -0.433 + 0.715LW$ ($R^2 = 0.987$). In each pear cultivar, ANN models were found to be more accurate in terms of both the training and testing phase than MLR models.

Research highlights: In the prediction of LA for different pear cultivars, ANN can thus be used in addition to MLR, as effective tools to circumvent difficulties met in the direct measurement of LA in the laboratory.

Additional keywords: *Pyrus communis* L.; artificial neural networks; multiple linear regressions; model estimation.

Abbreviations used: ANN (artificial neural network); L (length); LA (leaf area); LM (Levenberg Marquadt); MAD (mean absolute deviation); MAPE (mean absolute percentage error); MLR (multiple linear regression); MSE (mean square error); R^2 (determination coefficient); VIF (variance inflation factors); W (width).

Authors' contributions: Conceived, designed and performed the experiments: OA and DH. Analyzed the data: CB and KE. All authors wrote and approved the final manuscript.

Citation: Ozturk, A; Cemek, B; Demirsoy, H; Kucuktopcu, E (2019). Modelling of the leaf area for various pear cultivars using neuro computing approaches. Spanish Journal of Agricultural Research, Volume 17, Issue 4, e0206. <https://doi.org/10.5424/sjar/2019174-14675>

Received: 08 Feb 2019. **Accepted:** 11 Dec 2019.

Copyright © 2019 INIA. This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC-by 4.0) License.

Funding: The authors received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Correspondence should be addressed to Ahmet Öztürk: ozturka@omu.edu.tr

Introduction

The common pear (*Pyrus communis* L.) comes from the second largest temperate fruit tree globally and is cultivated worldwide. Overall, world pear production reached 24,168,309 tonnes in 2017, with Turkey producing 503,004 tonnes and thus ranking fifth place in pear production in the world that year (FAO, 2019). According to the FAO report on the state of world's plant genetic resources for food and agriculture at least 1,140 pear accessions are present in world-wide ex-situ collections (Federico *et al.*, 2008) and more than 600 cultivars have been reported in Turkey (Özbek, 1978).

Leaf area (LA) is commonly evaluated in fruit physiology experiments. The determination of LA is an important criterion in understanding respiration, transpiration, evapotranspiration, photosynthesis, light interception, water and nutrient use, fertilization, irrigation, plant growth, flowering, fruit set, yield, and fruit quality (Smith & Kliwer, 1984; Smart, 1985). The LA prediction models provide many benefits to researchers in horticultural experiments. Moreover, these models allow researchers to measure LA on the same plants during the plant growth period and may reduce variability in experiments (Gamiely *et al.*, 1991; NeSmith, 1992; Demirsoy, 2009).

The LA can be determined by both direct and indirect methods. The direct methods, called destructive methods, require destroying all the leaves in the plant canopy; therefore, it is not possible to take successive measurements from the same leaf (Fallovio *et al.*, 2008; Spann & Heerema, 2010). Direct methods are generally simple and accurate although the measurements involved can be laborious and time-consuming making it difficult to get a representative spatial sample and making large-scale implementation only marginally feasible (Rouphael *et al.*, 2006). Recently, new instruments, tools and machines such as hand-scanners and laser-optic devices have been developed for some LA measurements. However, these are too expensive and complex for basic and simple studies. In contrast, indirect methods, called non-destructive methods, are simple, rapid, inexpensive, and save labour relative to the direct measurements, and they can provide precise LA estimates without damaging the plant (Robbins & Pharr, 1987; Ahmadian-Moghadam, 2012).

Indirect methods are particularly useful if the empirical relations between LA and one or more dimensions of the leaf lamina need to be explained (Vazquez-Cruz *et al.*, 2012). The indirect method used most frequently necessitates estimating the LA with mathematical equations that involve linear measurements, including leaf length (L) and leaf width (W), or different combinations of these variables (Rouphael *et al.*, 2010; Giuffrida *et al.*, 2011; Walia & Kumar, 2017; Teobaldelli *et al.*, 2019). Numerous studies have been conducted with linear leaf measurements that have strong associations with LA in some fruit trees, such as banana (Potdar & Pawar, 1991), rabbiteye blueberry (NeSmith, 1991), white mulberry (Satpathy *et al.*, 1992), pecan (Sparks, 1996), blackberry (Uzun & Celik, 1999), peach (Demirsoy *et al.*, 2004), sweet cherry (Demirsoy & Demirsoy, 2003; Cittadini & Peri, 2006), strawberry (Demirsoy *et al.*, 2005), chestnut (Serdar & Demirsoy, 2006), kiwifruit (Mendoza-de Gyves *et al.*, 2007; Zenginbal *et al.*, 2007), hazelnut (Cristofori *et al.*, 2007), medlar (Mendoza-de Gyves *et al.*, 2008), persimmon (Cristofori *et al.*, 2008), coffee (Antunes *et al.*, 2008), vineyard (Tsialtas *et al.*, 2008; Sanchez-de Miguel *et al.*, 2011), citrus (Mazzini *et al.*, 2010), apple (Kishore *et al.*, 2012; Sala *et al.*, 2015), walnut (Keramatlou *et al.*, 2015), apricot (Cirillo *et al.*, 2017), cacao (Suarez Salazar *et al.*, 2018), or loquat (Teobaldelli *et al.*, 2019).

In recent years, a number of artificial neural network (ANN) studies have been performed in agricultural fields and successful results have been achieved. Yuan *et al.* (2017) evaluated the performance of random forest (RF), ANN, and support vector machine (SVM) regression models comparing them with a partial least-

squares regression (PLS) model for heterogeneous soybean crops. The results showed that the ANN model could be used for estimating leaf area index (LAI). Kumar *et al.* (2017) developed an ANN model including L and W as inputs and compared them with regression models. They found that the ANN model is more accurate than the regression model. Küçükönder *et al.* (2016) used ANN and regression analysis techniques for developing the best LA estimation model and concluded that the ANN approach can be used as an alternative method in estimating the LA. Shabani *et al.* (2017) used ANN in order to estimate the LA of different plants and stated that ANN provides a good estimation of LA. However, they emphasized that ANN is applicable for all plant species, but it is necessary to establish specific equations for each plant in the application of other models.

Though, in the literature, plenty of different equations have been derived for various plants and cultivars, the coefficients, as well as the kinds of equations, are plant specific (Kumar & Sharma, 2013). Although some studies including pear cultivars (*Pyrus pyrifolia* cv. 'Nijiseiki') were made to determine the LA in the shoots and spurs (Kumar *et al.*, 1977; Spann & Heerema, 2010), there have been no attempts to develop a non-destructive LA prediction model and ANN for only the pear. In the previous studies, other fruit species and pears were examined together. The development of the individual LA model for a single species is more important in terms of developing a basic and correct equation for the individual LA. Thus, this study aims to address this inadequacy by incorporating the ANN and comparing their performance for the pear by linear measurements of leaf geometry with multiple linear regressions (MLR) in pear.

Material and methods

The study was carried out on 12 pear cultivars in Samsun in Turkey from 2014 to 2015 to improve a leaf area prediction model and ANNs model. The examined pear cultivars were grafted on 'BA 29' quince rootstock. To develop the LA prediction model and validation, these 12 pear cultivars, which are commonly grown and economically important, with different numbers of leaves were selected (Table 1). The full expanded different sized leaf samples were randomly taken from the tree canopy in the actively growing season in one-month intervals (three months; June, July, August) during two investigation years, *i.e.* a total of 2975 leaves.

Initially, each leaf was taken and placed on an A4 sheet and copied (at a 1:1 ratio) with a photocopier.

Table 1. Descriptive statistics for pear cultivars used in this study.

Cultivars	Training				Testing			
	n	Length (cm)	Width (cm)	Leaf area (cm ²)	n	Length (cm)	Width (cm)	Leaf area (cm ²)
All cultivars	2975	6.669 ± 1.385	3.741 ± 1.109	18.240 ± 8.918	1262	6.932 ± 1.240	3.892 ± 0.893	19.221 ± 7.164
Abate Fetel	173	6.109 ± 0.882	3.265 ± 0.841	13.503 ± 5.412	74	6.738 ± 0.827	3.828 ± 0.780	17.208 ± 5.313
Akça	55	7.421 ± 0.636	5.118 ± 0.533	28.751 ± 5.400	22	7.345 ± 0.686	5.023 ± 0.581	27.950 ± 5.663
Carmen	317	6.425 ± 0.942	3.420 ± 0.619	16.010 ± 4.859	135	6.635 ± 0.993	3.552 ± 0.552	17.019 ± 4.979
Decana	248	5.990 ± 0.780	3.713 ± 0.663	16.050 ± 4.361	105	6.387 ± 0.802	3.777 ± 0.602	17.422 ± 4.279
Deveci	811	6.421 ± 1.507	3.172 ± 1.280	15.014 ± 10.157	346	7.101 ± 1.220	3.472 ± 0.781	17.170 ± 5.902
Etrusca	88	6.545 ± 0.740	3.535 ± 0.484	16.242 ± 3.960	37	6.516 ± 0.822	3.508 ± 0.571	16.081 ± 4.832
Kaiser Alexander	45	7.589 ± 0.913	4.944 ± 0.873	27.193 ± 7.526	18	7.217 ± 1.008	4.717 ± 1.115	24.917 ± 9.079
Limon	67	8.855 ± 1.271	5.140 ± 0.652	31.685 ± 7.293	28	8.882 ± 1.164	5.139 ± 0.527	31.611 ± 5.451
Kieffer	144	7.835 ± 1.100	4.712 ± 0.729	25.681 ± 7.291	60	8.038 ± 1.138	4.822 ± 0.768	27.015 ± 7.474
Samsun Güzeli	83	6.330 ± 0.963	3.801 ± 0.559	17.240 ± 4.865	35	6.140 ± 0.891	3.743 ± 0.534	16.549 ± 4.416
Santa Maria	376	7.607 ± 1.119	4.703 ± 0.937	25.553 ± 8.650	160	7.404 ± 1.072	4.562 ± 0.930	23.813 ± 8.019
Williams	568	6.375 ± 1.423	3.636 ± 0.746	16.903 ± 6.270	242	6.462 ± 1.338	3.849 ± 0.772	18.024 ± 6.365

A placom digital planimeter (Sokkisha Planimeter Inc., Model KP-90) was used to measure the actual leaf area of the copy. The leaf width (W) and length (L) of the leaf samples were also measured for model construction. W (cm) was measured from tip to tip at the widest part of the lamina and L (cm) was measured from the lamina tip to the point of petiole intersection along the midrib (Fig. 1). All values were recorded to the nearest 0.1 cm.

Model construction for multiple linear regressions (MLR)

The most common method used in the estimation of LA is to employ MLR. To determine the model, MLR

analysis using the stepwise method was employed. MLR analysis of the observed data was performed from 12 pear cultivars. For this reason, analysis was conducted with various subsets of the independent variables, namely, L, $L^{0.5}$, L^2 , W, $W^{0.5}$, W^2 , LW, $L^2 W^2$, (L+W) and $(L+W)^2$ to develop the best model for predicting LA by using the Microsoft Office 2015 Excel package program. The MLR analysis was carried out until the deviation sum of squares was minimized.

When two or more variables (*i.e.* L and W) in the model are correlated, the problem of collinearity may occur. The collinearity provides redundant information about the response. In order to detect collinearity, variance inflation factors (VIF) for each




Figure 1. Pear leaf showing the position of length (L) and width (W).

predictor were calculated (Mansfield & Helms, 1982). The VIF was calculated as, $VIF_j = 1/(1-R_j^2)$ where R_j^2 is the coefficient of determination of the model that includes all predictors except the j^{th} predictor. If VIF values are less than 10, then there is no problem with collinearity.

Artificial neural networks (ANN)

The multilayer perception (MLP) is the most broadly used model of neural networks (Fig. 2). The MLP comprises a set of simple interconnected units (neurons or nodes). The nodes are linked by weights and output signals which are a function of the sum of the inputs to the node modified by a simple nonlinear transfer, or activation, function (Haykin, 1994; Gardner & Dorling, 1998).

For processing ANN, the MATLAB (R2010b) software program was used. Levenberg Marquadt (LM) back propagation was employed to train the network as a learning algorithm. This algorithm was selected because it is a more effective and easy to learn model for complex networks than the standard back propagation (Sapna *et al.*, 2012). The ANN structure used in this study contains only one hidden layer, because many theoretical and experimental results confirm that one hidden layer can be enough for forecasting problems (Cybenko, 1989; Hornik *et al.*, 1989). Neurons in the input layer had no transfer function, while for hidden and output nodes there were tangent sigmoid (tansig) and linear transfer functions (purelin), respectively. The number of neurons in the hidden layer changed from 3 and 5 to 7 to achieve the optimal training network. To determine the final network architecture with the most

ideal neuron numbers, a trial-and-error procedure was undertaken, starting from 20 iteration steps to 100 steps in increases of 10 steps.

Data pre-processing

Data was divided into 70% for training and 30% for testing both MLR and ANN. For purposes of bringing all the data, that is, input and output data into a comparable range the training and testing data was standardized to fall in a range between 0 and 1 as given in formula (1) below,

$$X_n = \frac{X_i - X_{\min}}{X_{\max} - X_{\min}} \quad (1)$$

where X_n is the standardized value; X_i is the observed value; X_{\min} and X_{\max} are the minimum and maximum values of the training and testing data, respectively.

Standardization has the advantage of eliminating the effect of absolute magnitude values which neural networks are sensitive to. The performances of different models were evaluated by using the determination coefficient (R^2), mean square error (MSE), mean absolute deviation (MAD) and mean absolute percentage error (MAPE).

$$R^2 = \frac{\sum_{i=1}^n (LA_{mea} - \overline{LA}_{mea})(LA_{est} - \overline{LA}_{est})}{\sum_{i=1}^n (LA_{mea} - \overline{LA}_{mea})^2 \sum_{i=1}^n (LA_{est} - \overline{LA}_{est})^2} \quad (2)$$

$$MSE = \frac{\sum_{i=1}^n (LA_{mea} - LA_{est})^2}{n} \quad (3)$$




Figure 2. Architecture of the multilayer perception (MLP) applied to estimate the leaf area (LA).

$$MAD = \frac{\sum_{i=1}^n |LA_{mea} - LA_{est}|}{n} \quad (4)$$

$$MAPE = \frac{\sum_{i=1}^n \left| \frac{LA_{mea} - LA_{est}}{LA_{mea}} \right|}{n} \times 100 \quad (5)$$

where LA_{mea} is the measured LA value; LA_{est} is the estimated LA value; \overline{LA}_{mea} and \overline{LA}_{est} represent the average values of measured and estimated LA, respectively; and n is the number of data considered.

Higher values of R^2 and lower values of MSE, and MAD indicate a better forecast accurateness by the model. The R^2 measures the extent of the linear association of two variables and values range from 0 to 1. A value of 1 indicates a perfect agreement between the measured and predicted values and a value of 0 indicates no agreement which shows poor model performance. If MAPE values are less than 10%, the model is considered to have a high degree of accuracy (Lewis, 1982).

Results

MLR model

Collinearity between L and W was evaluated before the developed LA model in different pear varieties. As the VIF values were less than 10, there was no collinearity between L and W. Thus, these variables can be taken in the LA models. The best models of MLR for different pear cultivars are presented in Table 2. As seen in Table 2, both L and W variables are compulsory to estimate pear LA. Results showed that all MLR models provide a high degree of correlations between LA and L-W. The R^2 values were between 0.932 and 0.992, MAPE values were between 3.043% and 9.235%, MSE values were between 0.781 and 4.151, and MAD between 0.641 and 1.666 in all MLR models. Among these MLR models, the model for the 'Deveci' pear cultivar [$LA = -0.837 + 0.722LW$] was the best model based on the selection criteria (higher R^2 , and lowest

Table 2. Estimation performance of multiple linear regression (MLR) both training and testing phase.

Cultivars	Model form	Fitted coefficient and constant		Training				Testing			
		a	b	MAPE (%)	MSE (cm ²)	MAD (cm ²)	R ²	MAPE (%)	MSE (cm ²)	MAD (cm ²)	R ²
All cultivars	LA = a + bLW	-0.433 (0.044)	0.715 (0.001)	4.640	1.063	0.787	0.987	4.446	1.128	0.815	0.979
Abate Fetel	LA = a + b(L+W) ²	-1.636 (0.257)	0.167 (0.003)	7.150	1.229	0.883	0.958	5.924	1.490	0.975	0.949
Akça	LA = a + bL ² W ²	13.880 (0.418)	0.010 (0.001)	3.043	1.044	0.859	0.964	3.190	1.348	0.827	0.970
Carmen	LA = a + b(L+W)	-15.489 (0.437)	3.200 (0.044)	6.508	1.321	0.897	0.944	5.258	1.181	0.862	0.954
Decana	LA = a + bL ² W ²	8.062 (0.141)	0.015 (0.001)	5.447	1.071	0.796	0.944	4.391	0.966	0.751	0.948
Deveci	LA = a + bLW	-0.837 (0.058)	0.722 (0.002)	4.831	0.781	0.641	0.992	4.397	0.788	0.677	0.983
Etrusca	LA = a + bL ² W ²	8.200 (0.236)	0.014 (0.001)	4.957	0.893	0.751	0.943	6.133	2.072	0.840	0.932
Kaiser Alexander	LA = a + bL ² W ²	13.427 (0.590)	0.009 (0.001)	5.617	3.289	1.422	0.942	9.235	4.151	1.623	0.961
Limon	LA = a + bL ² W ²	15.641 (0.463)	0.007 (0.001)	3.933	2.303	1.666	0.957	3.763	2.184	1.169	0.935
Kieffer	LA = a + bL ² W ²	13.341 (0.234)	0.008 (0.001)	4.910	1.951	1.121	0.963	4.755	2.088	1.137	0.965
Samsun Güzeli	LA = a + bL ² W ²	8.384 (0.274)	0.014 (0.001)	6.288	1.351	0.954	0.943	6.359	1.178	0.862	0.945
Santa Maria	LA = a + bLW	-1.233 (0.195)	0.734 (0.005)	3.793	1.320	0.922	0.982	3.853	1.142	0.854	0.983
Williams	LA = a + b(L+W) ²	0.654 (0.166)	0.156 (0.002)	6.549	1.929	1.063	0.951	6.948	2.443	1.228	0.941

MAD: mean absolute deviation. MAPE: mean absolute percentage error. MSE: mean square error. R²: determination coefficient.

MAPE, MSE and MAD). Also, the general formula is obtained by using all pear cultivars [$LA = -0.433 + 0.715LW$]. Scatter plots for both training and testing for all pear cultivars and the 'Deveci' pear cultivar in the MLR model are shown in Fig. 3.

ANN model

The second approach to estimate LA is the ANN method. The best models of training and testing for ANN are presented in Table 3. According to the findings, MLP (2,3,1, LM) gives the best results for 'Deveci', with MAPE values of 4.293 and 3.426, MSE values of 0.524 and 0.530, MAD values of 0.539 and 0.500, and R^2 values of 0.995 and 0.986 for training and testing, respectively. This result means that the ANN model was able to explain 99.50% of variability in LA in training data, and 98.60% of variability in testing data, when L and W were used as input data. Results for the training and testing phase of all pear cultivars

and the 'Deveci' cultivar are also graphically given in Fig. 4. As can be seen in Table 3, 'Akça' also gives the worst estimate with the highest MSE and MAE. The weight and bias values used in the formulation of MLP (2,3,1, LM) are given in Tables 4 and 5.

Discussion

Leaf area is associated with many agronomical and physiological processes, including photosynthesis, respiration, transpiration, canopy light interception, irrigation, fertilization, pruning, fruit thinning and water use efficiency. As there is an integral relationship between leaves and many of these physiological processes, it is often necessary to estimate LA non-destructively in order not to disturb the system, or to allow for multiple measurements to be made. Because of the integral relationships between green leaves and the physiological processes detailed above, the excision




Figure 3. The relationship between measured and estimated leaf area (LA) for multiple linear regression (MLR) model. MAD: mean absolute deviation. MSE: mean square error. R^2 : determination coefficient.

Table 3. Estimation performance of ANN both training and testing phase.

Cultivars	Training				Testing			
	MAPE (%)	MSE (cm ²)	MAD (cm ²)	R ²	MAPE (%)	MSE (cm ²)	MAD (cm ²)	R ²
All cultivars	4.031	0.731	0.655	0.991	3.723	0.793	0.669	0.985
Abate Fetel	6.730	1.235	0.886	0.979	5.874	1.250	0.950	0.983
Akça	4.223	1.853	1.229	0.988	4.181	1.220	1.929	0.991
Carmen	4.824	0.866	0.755	0.977	4.191	0.719	0.823	0.979
Decana	3.328	0.453	0.514	0.977	3.093	0.532	0.500	0.978
Deveci	4.293	0.524	0.539	0.995	3.426	0.530	0.500	0.986
Etrusca	3.026	0.363	0.478	0.979	2.658	0.419	0.294	0.989
Kaiser Alexander	2.226	0.613	0.597	0.989	2.729	0.621	0.701	0.993
Limon	1.693	0.446	0.521	0.992	1.840	0.539	0.427	0.986
Kieffer	3.995	1.490	1.001	0.978	3.741	1.009	1.591	0.976
Samsun Güzeli	3.510	0.537	0.586	0.980	3.027	0.473	0.343	0.985
Santa Maria	3.181	0.905	0.744	0.989	3.834	0.815	1.091	0.987
Williams	3.914	0.633	0.615	0.986	3.535	0.593	0.633	0.985

MAD: mean absolute deviation. MAPE: mean absolute percentage error. MSE: mean square error. R²: determination coefficient.


**Figure 4.** The relationship between measured and estimated leaf area (LA) for artificial neural networks (ANN) model. MAD: mean absolute deviation. MSE: mean square error. R²: determination coefficient.

Table 4. The weight values between the input layer and the hidden layer.

Weights	Number of neurons in the hidden layer (i)		
	1	2	3
W_{1i}	-0.723	0.004	9.883
W_{2i}	-3.832	0.110	-2.503
Bias1	-35.733	92.017	17.104

of leaves from the plants is necessary. Furthermore, it is not possible to make successive measurements of the same leaf, and the plant canopy would be damaged and cause problems in other measurements (Tsialtas & Maslaris, 2005). This is why several combinations of measurements and models relating L and W to LA have been developed for several fruit trees; however, until now, no models have been developed for pears.

In this study, we developed a simple, easy and reliable LA prediction model for pear cultivars using MLR. There were no significant differences between the predicted and actual LA in any of the pear cultivars. For this reason, the model can be reliably used in pear physiological studies and research aiming to develop more productive and efficient pruning and training methods. In the previous studies, Kumar *et al.* (1977) and Spann & Heerema (2010) have developed a LA prediction model in some fruit species including pears.

Kumar *et al.* (1977) determined that the K factor in the pear was between 0.706 and 0.756 and showed that the formula can be used reliably. Spann & Heerema (2010), who tried to develop a LA prediction model in order to determine the total LA on a shoot for some fruit species, determined the R^2 of the developed model for the 'Nijiseiki' pear cultivar as 0.93. In our study, the confidence coefficient of the LA model was determined for both cultivars and in general was determined to be more reliable than in the previous studies. In addition, the size of the leaves on the plant may vary depending on the position of the leaf on the shoot or spur. Since the leaves on the spurs show a close spacing form, the necessity of determining their area individually is emphasized (Heerema *et al.*, 2008). The estimation model we obtained in this study was more effective than the previous studies on pears in predicting the area of the leaves from all different parts of the plant, and the study only included pears. In addition, validation of the models identified in this study was performed. Validation of the generated LA estimation models is very important for the reliability of the model (Demirsoy & Lang, 2010).

Confidence in the precision of these models would provide researchers with a reasonably fast and inexpensive method to use in studies on plant physiology such as respiration, transpiration and photosynthesis

without destructive leaf harvesting in different plant species such as sweet cherry (Demirsoy & Demirsoy, 2003), grape (Williams & Martinson, 2003; Tsialtas *et al.*, 2008), peach (Demirsoy *et al.*, 2004), strawberry (Mandal *et al.*, 2002; Demirsoy *et al.*, 2005), chestnut (Serdar & Demirsoy, 2006), hazelnut (Cristofori *et al.*, 2007), kiwi (Mendoza-de Gyves *et al.*, 2007; Zenginbal *et al.*, 2007), medlar (Mendoza-de Gyves *et al.*, 2008), persimmon (Cristofori *et al.*, 2008), pecan (Torri *et al.*, 2009), citrus (Mazzini *et al.*, 2010), pomegranate (Meshram *et al.*, 2012), walnut (Keramatlou *et al.*, 2015), and apricot (Cirillo *et al.*, 2017).

ANN applies for many objectives in agricultural research such as: crop yield and fruit weight prediction, evapotranspiration prediction, soil parameter estimation, water demand forecasting and hydrological forecasting (Shabani *et al.*, 2017). Therefore, ANNs are becoming a popular tool for modelling complex input-output dependencies (Maren *et al.*, 1990). Different researchers have shown that ANN models often give better results than traditional methods (Moosavi & Sepaskha, 2012). This ability of ANN models depends on modelling complexity and nonlinearity that is overlooked by traditional statistical regression models and is due to the architecture of an ANN, which allows highly correlated inputs to be used to enhance the ANN's modelling capability. Thus, ANN models are becoming a widespread prediction method because some statistical assumptions that are significant for forecasting regression models are now unnecessary (Ercanlı *et al.*, 2018).

Also, in addition to MLR, we developed ANN models to estimate LA. When comparing the results of two methods, ANN models were better than MLR with L, W inputs for LA estimation considering MSE, MAD, MAPE and R^2 criteria for both the training and testing phases (Tables 2 and 3). As the ANN models use non-linear relations with input and output data, we were able to estimate LA with a high degree of accuracy as compared to the MLR. In the prediction of LA for different pear cultivars, ANNs can thus be used in addition to MLR, as an effective tool to circumvent difficulties in the direct measurement of LA in the laboratory. Similar results were obtained by Ahmadian-Moghadam (2012), Küçükönder *et al.* (2016), and Kumar *et al.* (2017). From these results, it can be concluded that ANN can be used for all plants, but a

Table 5. The weight values between the hidden layer and the output layer.

Weights	Number of neurons in the hidden layer (i)			
	1	2	3	Bias2
W_i	-6.570	-1.548	-3.993	27.785

specific equation should be prepared in applications of other methods for each plant.

In conclusion, in this study, different learning algorithms of ANNs and MLR were employed and their performances assessed by using L and W for estimating LA. The results of ANNs were compared to those of MLR and one another statistically using R^2 , MSE and MAD and MAPE. Comparative analysis of the network model results indicated that all models generally performed well for LA estimation during training and testing phases. It is clear from the study that the general predictions by all ANNs and MLR models were good, with ANN models being slightly better in predicting LA than MLR. As a result of this, in the prediction of LAs for different pear cultivars, ANNs can thus be used in addition to MLR, as an effective tool to circumvent difficulties met in the direct measurement of LAs in the laboratory.

References

- Ahmadian-Moghadam H, 2012. Prediction of pepper (*Capsicum annuum* L.) leaf area using group method of data handling-type neural networks. *Int J Agr Sci* 2: 993-999.
- Antunes WC, Pompelli MF, Carretero DM, DaMatta FM, 2008. Allometric models for non-destructive leaf area estimation in coffee (*Coffea arabica* and *Coffea canephora*). *Ann Appl Biol* 153: 23-40. <https://doi.org/10.1111/j.1744-7348.2008.00235.x>
- Cirillo C, Pannico A, Basile B, Rivera C, Giaccone M, Colla G, De Pascale S, Roupheal Y, 2017. A simple and accurate allometric model to predict single leaf area of twenty-one European apricot cultivars. *Eur J Hort Sci* 82: 65-71.
- Cittadini ED, Peri PL, 2006. Estimation of leaf area in sweet cherry using a non-destructive method. *Rev Invest Agropec* 35 (1): 143-150. <http://agris.fao.org/agris-search/search.do?recordID=NL2012024106>
- Cristofori V, Roupheal Y, Mendoza-de Gyves E, Bignami C, 2007. A simple model for estimating leaf area of hazelnut from linear measurements. *Sci Hort* 113: 221-225. <https://doi.org/10.1016/j.scienta.2007.02.006>
- Cristofori V, Fallovo C, Mendoza-de Gyves E, Rivera CM, Bignami C, Roupheal Y, 2008. Non-destructive, analogue model for leaf area estimation in Persimmon (*Diospyros kaki* L.F.) based on leaf length and width measurement. *Eur J Hort Sci* 73: 216-221.
- Cybenko G, 1989. Approximation by superpositions of a sigmoidal function. *Math Control Signals Syst* 2: 303-314. <https://doi.org/10.1007/BF02551274>
- Demirsoy H, Demirsoy L, 2003. Leaf area estimation model for some local cherry genotypes in Turkey. *Pak J Bot* 6 (2): 153-156. <https://doi.org/10.3923/pjbs.2003.153.156>
- Demirsoy H, Demirsoy L, Uzun S, Ersoy B, 2004. Non-destructive leaf area estimation in peach. *Eur J Hort Sci* 69: 144-146. <https://doi.org/10.1051/fruits:2005014>
- Demirsoy H, Demirsoy L, Öztürk A, 2005. Improved model for the non-destructive estimation of strawberry leaf area. *Fruits* 60: 69-73. <https://doi.org/10.1051/fruits:2005014>
- Demirsoy H, 2009. Leaf area estimation in some species of fruit tree by using models as a non-destructive method. *Fruits* 64: 45-51. <https://doi.org/10.1051/fruits/2008049>
- Demirsoy H, Lang GA, 2010. Validation of a leaf area estimation model for sweet cherry. *Span J Agric Res* 8 (3): 830-832. <https://doi.org/10.5424/sjar/2010083-1285>
- Ercanlı İ, Günlü A, Şenyurt M, Keleş S, 2018. Artificial neural network models predicting the leaf area index: a case study in pure even-aged Crimean pine forests from Turkey. *For Ecosyst* 5 (29): 1-12. <https://doi.org/10.1186/s40663-018-0149-8>
- Fallovo C, Cristofori V, de-Gyves EM, Rivera CM, Rea R, Fanasca S, Bignami C, Sassine Y, Roupheal Y, 2008. Leaf area estimation model for small fruits from linear measurements. *HortScience* 43: 2263-2267. <https://doi.org/10.21273/HORTSCI.43.7.2263>
- FAO, 2019. FAO Statistics database. <http://www.fao.org/faostat/en/#home>
- Federico M, Busconi M, Camangi F, Fogher C, Stefani A, Sebastiani L, 2008. Ancient Pomoideae (*Malus domestica* Borkh. and *Pyrus communis* L.) cultivars in "Appennino Toscano" (Tuscany, Italy): molecular (SSR) and morphological characterization. *Caryologia* 61: 320-331. <https://doi.org/10.1080/00087114.2008.10589643>
- Gamiely S, Randle WM, Mills HA, Smittle DA, 1991. A rapid and non-destructive method for estimating leaf area of onions. *HortScience* 26 (2): 206. <https://doi.org/10.21273/HORTSCI.26.2.206>
- Gardner MW, Dorling SR, 1998. Artificial neural networks (the multilayer perceptron) - A review of applications in the atmospheric sciences. *Atmos Environ* 32: 2627-2636. [https://doi.org/10.1016/S1352-2310\(97\)00447-0](https://doi.org/10.1016/S1352-2310(97)00447-0)
- Giuffrida F, Roupheal Y, Toscano S, Scuderi D, Romano D, Rivera C, Colla G, Leonardi C, 2011. Simple model for nondestructive leaf area estimation in bedding plants. *Photosynthetica* 49: 380-388. <https://doi.org/10.1007/s11099-011-0041-z>
- Haykin S, 1994. Neural networks: a comprehensive foundation. Prentice Hall PTR.
- Heerema RJ, Weinbaum SA, Pernice F, DeJong TM, 2008. Spur survival and return bloom in almond [*Prunus dulcis* (Mill.) D.A. Webb] varied with spur fruit load, specific leaf weight, and leaf area. *J Hort Sci Biotechnol* 83: 274-281. <https://doi.org/10.1080/14620316.2008.11512380>
- Hornik K, Stinchcombe M, White H, 1989. Multilayer feedforward networks are universal approximators. *Neural*

- Networks 2: 359-366. [https://doi.org/10.1016/0893-6080\(89\)90020-8](https://doi.org/10.1016/0893-6080(89)90020-8)
- Keramatlou I, Sharifani M, Sabouri H, Alizadeh M, Kamkar B, 2015. A simple linear model for leaf area estimation in Persian walnut (*Juglans regia* L.). *Sci Hort* 184: 36-39. <https://doi.org/10.1016/j.scienta.2014.12.017>
- Kishore D, Pramanick K, Verma J, Singh R, 2012. Non-destructive estimation of apple (*Malus domestica* Borkh.) leaf area. *J Hort Sci Biotechnol* 87: 388-390. <https://doi.org/10.1080/14620316.2012.11512881>
- Küçükönder H, Boyacı S, Akyüz A, 2016. A modeling study with an artificial neural network: developing estimation models for the tomato plant leaf area. *Turk J Agric For* 40: 203-212. <https://doi.org/10.3906/tar-1408-28>
- Kumar K, Srivastava RP, Singh AK, Bana DS, 1977. Use of linear measurement in the estimation of leaf area of some apricot, peach, plum, pear and guava varieties. *Ind J Hort* 34: 229-234.
- Kumar R, Sharma S, 2013. Simulation and validation of leaf area prediction model for *Picrorhiza kurroa* an endangered medicinal plant of Western Himalaya. *J Med Plants Res* 7: 1467-1473.
- Kumar K, Kumar S, Sankar V, Sakthivel T, Karunakaran G, Tripathi P, 2017. Non-destructive estimation of leaf area of durian (*Durio zibethinus*)-An artificial neural network approach. *Sci Hort* 219: 319-325. <https://doi.org/10.1016/j.scienta.2017.03.028>
- Lewis CD, 1982. Industrial and business forecasting methods: A practical guide to exponential smoothing and curve fitting. Butterworth-Heinemann.
- Mandal KK, Ghosh SK, Gayen P, 2002. A non-destructive way of leaf area estimation in the strawberry. *Ann Biol* 18 (1): 19-24.
- Mansfield ER, Helms BP, 1982. Detecting multicollinearity. *The Am Statist* 36: 158-160. <https://doi.org/10.1080/00031305.1982.10482818>
- Maren AJ, Harston CT, Pap RM, 1990. Handbook of neural computing applications. Acad Press, San Diego (CA), USA.
- Mazzini RB, Ribeiro RV, Pio RM, 2010. A simple and non-destructive model for individual leaf area estimation in citrus. *Fruits* 65: 269-275. <https://doi.org/10.1051/fruits/2010022>
- Mendoza-de Gyves E, Roupael Y, Cristofori V, Rosana Mira F, 2007. A non-destructive, simple and accurate model for estimating the individual leaf area of kiwi (*Actinidia deliciosa*). *Fruits* 62: 171-176. <https://doi.org/10.1051/fruits:2007012>
- Mendoza-de Gyves E, Cristofori V, Fallovo C, Roupael Y, Bignami C, 2008. Accurate and rapid technique for leaf area measurement in medlar (*Mespilus germanica* L.). *Adv Hort Sci* 22 (3): 223-226.
- Meshram DT, Gorantiwar SD, Singh NV, Suroshe SS, 2012. Non-destructive leaf area estimation in pomegranate cv: Bhagwa (*Punica granatum* L.). *Ind J Hort* 69: 163-167.
- Moosavi AA, Sepaskha A, 2012. Artificial neural networks for predicting unsaturated soil hydraulic characteristics at different applied tensions. *Arch Agron Soil Sci* 58 (2): 125-153. <https://doi.org/10.1080/03650340.2010.512289>
- NeSmith DS, 1991. Non-destructive leaf area estimation of rabbiteye blueberries. *HortScience* 26: 132. <https://doi.org/10.21273/HORTSCI.26.10.1332>
- NeSmith DS, 1992. Estimating summer squash leaf area nondestructively. *HortScience* 27: 77. <https://doi.org/10.21273/HORTSCI.27.1.77>
- Özbek S, 1978. Özel meyvecilik. Çukurova Üniversitesi Ziraat Fakültesi Yayınları (Turkish) 128: 392-483.
- Potdar M, Pawar K, 1991. Non-destructive leaf area estimation in banana. *Sci Hort* 45: 251-254. [https://doi.org/10.1016/0304-4238\(91\)90070-F](https://doi.org/10.1016/0304-4238(91)90070-F)
- Robbins NS, Pharr DM, 1987. Leaf area prediction models for cucumber from linear measurements. *HortScience* 22 (6): 1264-1266.
- Roupael Y, Rivera CM, Cardarelli M, Fanasca S, Colla G, 2006. Leaf area estimation from linear measurements in zucchini plants of different ages. *J Hort Sci Biotechnol* 81: 238-241. <https://doi.org/10.1080/14620316.2006.11512056>
- Roupael Y, Mouneimne A, Ismail A, Mendoza-De Gyves E, Rivera C, Colla G, 2010. Modeling individual leaf area of rose (*Rosa hybrida* L.) based on leaf length and width measurement. *Photosynthetica* 48: 9-15. <https://doi.org/10.1007/s11099-010-0003-x>
- Sala F, Arsene GG, Iordănescu O, Boldea M, 2015. Leaf area constant model in optimizing foliar area measurement in plants: A case study in apple tree. *Sci Hort* 193: 218-224. <https://doi.org/10.1016/j.scienta.2015.07.008>
- Sanchez-de-Miguel P, Junquera P, de la Fuente M, Jimenez L, Linares R, Baeza P, Lissarrague JR, 2011. Estimation of vineyard leaf area by linear regression. *Span J Agric Res* 9 (1): 202-212. <https://doi.org/10.5424/sjar/20110901-354-10>
- Sapna S, Tamilarasi A, Kumar MP, 2012. Backpropagation learning algorithm based on Levenberg Marquardt Algorithm. *Comp Sci Inform Technol (CS and IT)* 2: 393-398. <https://doi.org/10.5121/csit.2012.2438>
- Satpathy B, Shivnath A, Rao KM, Ghosh PL, Nair BP, 1992. An easy and rapid method of leaf area estimation in white mulberry (*Morus alba*). *Ind J Agric Sci* 62: 489-491.
- Serdar U, Demirsoy H, 2006. Non-destructive leaf area estimation in chestnut. *Sci Hort* 108 (2): 227-230. <https://doi.org/10.1016/j.scienta.2006.01.025>
- Shabani A, Ghaffary KA, Sepaskhah AR, Kamgar-Haghighi AA, 2017. Using the artificial neural network to estimate leaf area. *Sci Hort* 216: 103-110. <https://doi.org/10.1016/j.scienta.2016.12.032>

- Smart RE, 1985. Principles of grapevine canopy microclimate manipulation with implications for yield and quality: A review. *Am J Enol Viticul* 36: 230-239.
- Smith R, Kliewer W, 1984. Estimation of Thompson Seedless grapevine leaf area. *Am J Enol Viticult* 35: 16-22.
- Spann TM, Heerema RJ, 2010. A simple method for non-destructive estimation of total shoot leaf area in tree fruit crops. *Sci Hortic* 125: 528-533. <https://doi.org/10.1016/j.scienta.2010.04.033>
- Sparks D, 1996. A rapid method for estimating the leaf area of the Stuart pecan, *Carya illinoensis* Koch. *HortScience* 1: 93-94.
- Suarez Salazar J, Melgarejo L, Duran Bautista E, Di Rienzo J, Casanoves F, 2018. Non-destructive estimation of the leaf weight and leaf area in cacao (*Theobroma cacao* L.). *Sci Hortic* 229: 19-24. <https://doi.org/10.1016/j.scienta.2017.10.034>
- Teobaldelli M, Roupael Y, Fascelle G, Cristofori V, Rivera CM, Basile B, 2019. Developing an accurate and fast non-destructive single leaf area model for loquat (*Eriobotrya japonica* Lindl) cultivars. *Plants* 8 (7): 230-242. <https://doi.org/10.3390/plants8070230>
- Torri SI, Descalzi C, Frusso E., 2009. Estimation of leaf area in pecan cultivars (*Carya illinoensis*). *Cienc Investig Agrar* 36: 53-58. <https://doi.org/10.4067/S0718-16202009000100004>
- Tsialtas JT, Maslaris N, 2005. Leaf area estimation in a sugar beet cultivar by linear models. *Photosynthetica*, 43 (3): 477-479. <https://doi.org/10.1007/s11099-005-0077-z>
- Tsialtas JT, Koundouras S, Zioziou E, 2008. Leaf area estimation by simple measurements and evaluation of leaf area prediction models in Cabernet-Sauvignon grapevine leaves. *Photosynthetica* 46: 452-456. <https://doi.org/10.1007/s11099-008-0077-x>
- Uzun S, Çelik H, 1999. Leaf area prediction models (uzçelik-1) for different horticultural plants. *Turk J Agric For* 23: 645-650.
- Vazquez-Cruz M, Luna-Rubio R, Contreras-Medina L, Torres-Pacheco I, Guevara-Gonzalez R, 2012. Estimating the response of tomato (*Solanum lycopersicum*) leaf area to changes in climate and salicylic acid applications by means of artificial neural networks. *Biosyst Eng* 112: 319-327. <https://doi.org/10.1016/j.biosystemseng.2012.05.003>
- Walia S, Kumar R, 2017. Development of the nondestructive leaf area estimation model for valeriana (*Valeriana jatamansi* Jones). *Commun Soil Sci Plan* 48: 83-91. <https://doi.org/10.1080/00103624.2016.1253723>
- Williams L, Martinson TE, 2003. Non-destructive leaf area estimation of 'Niagara' and 'DeChaunac' grapevines. *Sci Hortic* 98: 493-498. [https://doi.org/10.1016/S0304-4238\(03\)00020-7](https://doi.org/10.1016/S0304-4238(03)00020-7)
- Yuan H, Yang G, Li C, Wang Y, Liu J, Yu H, Feng H, Xu B, Zhao X, Yang X, 2017. Retrieving soybean leaf area index from unmanned aerial vehicle hyperspectral remote sensing: Analysis of RF, ANN, and SVM regression models. *Remote Sens* 9: 309. <https://doi.org/10.3390/rs9040309>
- Zenginbal H, Uzun S, Ozcan M, Demir T, 2007. Leaf area prediction model for some kiwifruit (*Actinidia chinensis* Planch.) cultivars. *J Plant Sci* 2: 470-474. <https://doi.org/10.3923/jps.2007.470.474>