

Adnan S, Maltamo M, Coomes DA, Valbuena R, 2017. Effects of Plot Size, Stand Density and Scan Density on the Relationship between Airborne Laser Scanning Metrics and the Gini Coefficient of Tree Size Inequality. *Can J Forest Res* 47 (12): 1590-1602. <https://doi.org/10.1139/cjfr-2017-0084>

Adnan S, Maltamo M, Coomes DA, García-Abril A, Malhi Y, Manzanera JA, Butt N, Morecroft M, Valbuena R, 2018. A simple approach to forest structure classification using airborne laser scanning that can be adopted across bioregions. *Forest Ecol Manag* 433: 111-121. <https://doi.org/10.1016/j.foreco.2018.10.057>

Alberdi I, Vallejo R, Álvarez-González JG, Condés S, González-Ferreiro E, Guerrero S, Hernández L, Martínez-Jáuregui M, Montes F, Oliveira N, et al., 2017. The multi-objective Spanish National Forest Inventory. *Forest Syst* 26 (2): e04S. <https://doi.org/10.5424/fs/2017262-10577>

Alonso-Benito A, Arroyo LA, Arbelo M, Hernández-Leal P, 2016. Fusion of WorldView-2 and LiDAR data to map fuel types in the Canary Islands. *Remote Sens-Basel* 8 (8): 669. <https://doi.org/10.3390/rs8080669>

Álvarez-Martínez JM, Jiménez-Alfaro B, Barquín J, Ondiviela B, Recio M, Silió-Calzada A, Juanes JA, 2017. Modelling the area of occupancy of habitat types with remote sensing. *Methods Ecol Evol* 9: 580-593. <https://doi.org/10.1111/2041-210X.12925>

Álvarez-Taboada MF, Cimadevila HL, Rodríguez Pérez JR, Picos Martín J, 2004. Workflow to improve the forest management of *Eucalyptus globulus* stands affected by *Gonipterus scutellatus* in Galicia, Spain using remote sensing and GIS, Proc. SPIE 5574, Remote Sensing for Environmental Monitoring, GIS Applications, and Geology IV, (22 October 2004).

Álvarez-Taboada MF, 2006. Remote sensing and Geoinformation systems applied to the forest management of *Eucalyptus globulus* Labill. Stands damaged by *Gonipterus scutellatus* Gyllendall in Galicia. Doctoral Thesis. Universidade de Vigo. 319 pp.

Álvarez-Taboada MF, Lorenzo Cimadevila H, Wulder M, 2007a. Monitorización del estado sanitario de las masas de *Eucalyptus globulus* en Galicia empleando modelos de proceso, SIG y teledetección. Proc 2º simposio iberoamericano de Eucalipto Globulus in Vigo (Spain). October 17-20, CIDEU 4 vol II, pp 41-47.

Álvarez-Taboada MF, Rodríguez-Pérez JR, Castedo-Dorado F, Vega-Nieva D, 2007b. An operational protocol for post-fire evaluation at landscape scale in an object-oriented environment. Proceedings of the 6th International Workshop of the EARSeL Special Interest Group on Forest Fires JRC 8072. pp 202 – 207 (2007). 6th International Workshop of the EARSeL Special Interest Group on Forest Fires. Tesalonica, Grecia.

Álvarez-Taboada, F, Sanz-Ablanedo, E, Rodríguez Pérez, JR, Castedo-Dorado F, Lombardero MJ, 2014. Multi-sensor and multi-scale system for monitoring forest health in *Pinus radiata* stands defoliated by *Lymantria dispar* in NW Spain. Proceedings of the ForestSAT Open Conference System, <http://ocs.agr.unifi.it/index.php/forestsat2014/ForestSAT2014/paper/view/245>

Aragónés D, Rodríguez-Galiano V, Caparros-Santiago JA, Navarro-Cerrillo RM, 2017. El uso de la fenología de la superficie terrestre para discriminar entre especies de pinos mediterráneos. Nuevas plataformas y sensores de teledetección, XVII Congreso de la Asociación Española de Teledetección (Eds. Ruiz LA, Estornell J, Erena M), Murcia (Spain), October 3-7, pp: 219-222.

Arellano S, Vega JA, Rodríguez y Silva F, Fernández C, Vega-Nieva D, Álvarez-González JG, Ruiz-González AD, 2017. Validación de los índices de teledetección dNBR y RdNBR para determinar la severidad del fuego en el incendio forestal de Oia-O Rosal (Pontevedra) en 2013. *Revista de Teledetección* 49: 49-61. <https://doi.org/10.4995/raet.2017.7137>

Arellano-Pérez S, Castedo-Dorado F, López-Sánchez C, González-Ferreiro E, Yang Z, Díaz-Varela R, Ruiz-González A, 2018. Potential of Sentinel-2A Data to Model Surface and Canopy Fuel Characteristics in Relation to Crown Fire Hazard. *Remote Sens-Basel* 10(10): 1645. <https://doi.org/10.3390/rs10101645>

Arias-Rodil M, Diéguez-Aranda U, Álvarez-González JG, Pérez-Cruzado C, Castedo-Dorado F, González-Ferreiro E, 2018. Modeling diameter distributions in radiata pine plantations in Spain with existing countrywide LiDAR data. *Ann For Sci* 75 (2): 1-12.

Arozarena A, Villa G, Hermosilla J, Papí F, Valcárcel N, Peces JJ, Doménech E, García C, Tejeiro JA, 2006. El Plan Nacional de Observación del Territorio en España: situación actual y próximos pasos. *Mapping Interactivo* 111: 16-22.

Arroyo LA, Healey SP, Cohen WB, Cocero D, Manzanera JA, 2006. Using object-oriented classification and high-resolution imagery to map fuel types in a Mediterranean region. *J Geophys Res* 111: G04S04. <https://doi.org/10.1029/2005JG000120>

Askne JIH, Santoro M, Smith G, Fransson JES, 2003. Multitemporal Repeat-Pass SAR Interferometry of Boreal Forests. *IEEE Trans Geosci Rem Sens* 41: 1540-1550. <https://doi.org/10.1109/TGRS.2003.813397>

Aulló-Maestro I, Gómez C, Cuevas R, Rubio A, Montes F, 2017. Dinámica forestal de *Pinus sylvestris* L. y *Quercus pyrenaica* Willd. en el bosque de Hoyocasero (Ávila) mediante serie temporal Landsat (1984-2016) y métodos geoestadísticos. *Nuevas plataformas y sensores de teledetección, XVII Congreso de la Asociación Española de Teledetección* (Eds. Ruiz LA, Estornell J, Erena M), Murcia (Spain), October 3-7, pp: 143-146.

Axelsson C, Skidmore AK, Schlerf M, Fauzi A, Verhoef W, 2012. Hyperspectral analysis of mangrove foliar chemistry using PLSR and support vector regression. *Int J Remote Sens* 34: 1724-1743. <https://doi.org/10.1080/01431161.2012.725958>

Banskota A, Kayastha N, Falkowski MJ, Wulder MA, Froese RE, White JC, 2014. Forest Monitoring Using Landsat Time Series Data: A Review. *Can J Remote Sens* 40: 362-384. <https://doi.org/10.1080/07038992.2014.987376>

Bastarrika A, Alvarado M, Artano K, Martínez MP, Mesanza A, Torre L, Ramo R, Chuvieco E, 2014. BAMS: A Tool for Supervised Burned Area Mapping Using Landsat Data. *Remote Sens-Basel* 6: 12360-12380. <https://doi.org/10.3390/rs61212360>

Belenguer-Plomer MA, Tanase MA, Fernández-Carrillo A, Chuvieco E, 2018. Insights into burned areas detection from Sentinel-1 data and locally adaptive algorithms, *Proc. SPIE 10790, Earth Resources and Environmental Remote Sensing/GIS Applications IX*, 107901S (9 October 2018).

Belward AS, Skøien JO, 2015. Who launched what, when and why; trends in global land-cover observation capacity from civilian earth observation satellites. *ISPRS J Photogramm*103: 115-128.

Bengoa JL, De Blanco V, Nafria DA, 2017. Clasificación semiautomática de cubiertas naturales

arboladas en Castilla y León. 7º Congreso Forestal Español. 26-30 de junio de 2017. Plasencia, Cáceres, España.

Blanco-Martínez J, Rodríguez F, Martínez S, Martínez AA, García JB, Fernández JJ, Roldán A, Diez FJ, Lizarralde I, Cabrera M, 2017. Generación de un inventario forestal regional y una cartografía de modelos de combustible para Castilla-La Mancha. 7 Congreso Forestal Español, 26-30 de junio, Plasencia, Spain.

Blázquez-Casado Á, González-Olabarria JR, Martín-Alcón S, Just A, Cabré M, Coll LI, 2015. Assessing post-storm forest dynamics in the Pyrenees using high-resolution LiDAR data and aerial photographs. *J Mt Sci* 12 (4): 841. <https://doi.org/10.1007/s11629-014-3327-3>

Bonal R, Vargas-Osuna E, Mena JD, Aparicio JM, Santoro M, Martín A, 2018. Looking for variable molecular markers in the chestnut gall wasp *Dryocosmus kuriphilus*: First comparison across genes *Scientific Reports*. 8. 10.1038/s41598-018-23754-z. <https://doi.org/10.1038/s41598-018-23754-z>

Borràs J, Delegido J, Pezzola A, Pereira M, Morassi G, Camps-Valls G, 2017. Clasificación de usos del suelo a partir de imágenes Sentinel-2. *Revista de Teledetección* 48: 55-66. <https://doi.org/10.4995/raet.2017.7133>

Botella-Martínez MA, Fernández-Manso A, 2017. Estudio de la severidad post-incendio en la Comunidad Valenciana comparando los índices dNBR, RdNBR y RBR a partir de imágenes Landsat 8. *Revista de Teledetección* 49: 33-47. <https://doi.org/10.4995/raet.2017.7095>

Bottalico F, Chirici G, Giannini R, Mele S, Mura M, Puxeddu M, McRoberts RE, Valbuena R, Travaglini D, 2017. Modeling Mediterranean Forest Structure Using Airborne Laser Scanning Data. *Int J Appl Earth Obs* 57: 145-153. <https://doi.org/10.1016/j.jag.2016.12.013>

Bradley BA, 2014. Remote Detection of Invasive Plants: A Review of Spectral, Textural and Phenological Approaches. *Biol Invasions* 16: 1411-1425. <https://doi.org/10.1007/s10530-013-0578-9>

Cabello J, Alcaraz-Segura D, Reyes A, Lourenço P, Requena JM, Bonache J, Castillo P, Valencia S, Naya J, Ramírez L, Serrada J, 2016. Sistema para el seguimiento del funcionamiento de ecosistemas en la Red de Parques Nacionales de España mediante teledetección. *Revista de Teledetección* 46: 119-131. <https://doi.org/10.4995/raet.2016.5731>

Camarero JJ, Bigler C, Linares JC, Gil-Pelegrín E, 2011. Synergistic effects of past historical logging and drought on the decline of Pyrenean silver fir forests. *For Ecol Manage* 262: 759-769.

Cardil A, Vepakomma U, Brotons LI, 2017. Assessing processionary moth defoliation using unmanned aerial systems. *Forests* 8: 402. <https://doi.org/10.3390/f8100402>

Carter G, 1993. Responses of leaf spectral reflectance to plant stress. *Am J Bot* 80: 231-243. <https://doi.org/10.1002/j.1537-2197.1993.tb13796.x>

Cartus O, Santoro, M., Kelldorfer J, 2012. Mapping forest aboveground biomass in the Northeastern United States with ALOS PALSAR dual-polarization L-band. *Remote Sens Environ* 124: 466-478. <https://doi.org/10.1016/j.rse.2012.05.029>

Castedo-Dorado F, Lago-Parra G, Lombardero MJ, Liebhold AM, Álvarez-Taboada F, 2016. European gypsy moth (*Lymantria dispar dispar* L.) completes development and defoliates exotic radiata pine

- plantations in Spain. *New Zeal J For Sci* 46: 18. <https://doi.org/10.1186/s40490-016-0074-y>
- Chrysafis I, Mallinis G, Gitas I, Tsakiri-Strati M, 2017. Estimating Mediterranean forest parameters using multi seasonal Landsat 8 OLI imagery and an ensemble learning method. *Remote Sens Environ* 199: 154-166. <https://doi.org/10.1016/j.rse.2017.07.018>
- Chuvienco E, Riaño D, Aguado I, Cocero D, 2002. Estimation of fuel moisture content from multitemporal analysis of Landsat Thematic Mapper reflectance data: applications in fire danger assessment. *Int J Remote Sens* 23 (11): 2145-2162. <https://doi.org/10.1080/01431160110069818>
- Chuvienco E, Aguado I, Cocero D, Riaño D, 2003. Design of an empirical index to estimate fuel moisture content from NOAA-AVHRR analysis in forest fire danger studies. *Int J Remote Sens* 24(8): 1621-1637. <https://doi.org/10.1080/01431160210144660b>
- Chuvienco E, Cocero D, Riaño D, Martínez P, Martínez-Vega J, De la Riva J, Pérez F, 2004a. Combining NDVI and surface temperature for the estimation of live fuel moisture content in forest fire danger rating. *Remote Sens Environ* 92: 322-331. <https://doi.org/10.1016/j.rse.2004.01.019>
- Chuvienco E, Cocero D, Aguado I, Palacios-Orueta A, Prado E, 2004b. Improving burning efficiency estimates through satellite assessment of fuel moisture content. *J Geoph Res-Atmos* 109: D14S07.
- Chuvienco E, De Santis A, Riaño D, Halligan K, 2007. Simulation approaches for burn severity estimation using remotely sensed images. *Fire Eco* 3 (1): 129-150. <https://doi.org/10.4996/fireecology.0301129>
- Cicuéndez V, Litago J, Huesca M, Rodríguez-Rastrero M, Recuero L, Merino de Miguel S, Palacios-Orueta A, 2015. Assessment of the gross primary production dynamics of a Mediterranean holm oak forest by remote sensing time series analysis. *Agroforest Syst* 89 (3): 491-510. <https://doi.org/10.1007/s10457-015-9786-x>
- Cifuentes JM, Fernández-Manso A, Sanz-Ablanedo E, 2017. Utilización de vehículo aéreo no tripulado (VANT) en el estudio de los niveles de severidad por chancro del castaño en el NO de España. In: *Nuevas plataformas y sensores de teledetección. Nuevas plataformas y sensores de teledetección, XVII Congreso de la Asociación Española de Teledetección* (Eds. Ruiz LA, Estornell J, Erena M), Murcia (Spain), October 3-7, pp: 477-480.
- Clark ML, Roberts DA, Clark DB, 2005. Hyperspectral discrimination of tropical rain forest tree species at leaf to crown scales. *Remote Sens Environ* 96: 375-398. <https://doi.org/10.1016/j.rse.2005.03.009>
- Claverie M, Ju J, Masek JG, Dungan JL, Vermote EF, Roger J-C, Skakun SV, Justice C, 2018. The harmonized Landsat and Sentinel-2 surface reflectance data set. *Remote Sens Environ* 219: 145-161. <https://doi.org/10.1016/j.rse.2018.09.002>
- Cloude SR, Papathanassiou KP, 1998. Polarimetric SAR Interferometry. *IEEE Trans Geosci Rem Sens* 36: 1551-1565. <https://doi.org/10.1109/36.718859>
- Cohen WB, Goward SN, 2004. Landsat's role in ecological applications of remote sensing. *Biosciences* 54 (6): 535-545. [https://doi.org/10.1641/0006-3568\(2004\)054\[0535:LRIEAO\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2004)054[0535:LRIEAO]2.0.CO;2)
- Cohen WB, Yang Z, Stehman SV, Schroeder TA, Bell DM, Masek JG, Huang Ch, Meighs GW, 2016.

Forest disturbance across the conterminous United States from 1985-2012: the emerging dominance of forest decline. *Forest Ecol Manag* 360: 242-252. <https://doi.org/10.1016/j.foreco.2015.10.042>

Cohen WB, Yang Z, Healey SP, Kennedy RE, Gorelick N, 2018. A LandTrendr multispectral ensemble for forest disturbance detection. *Remote Sens Environ* 205: 131-140. <https://doi.org/10.1016/j.rse.2017.11.015>

Condés S, Fernández-Landa A, Rodríguez F, 2013. Influencia del inventario de campo en el error de muestreo obtenido en un inventario con tecnología LiDAR. 6º Congreso Forestal Español. SECF

Crespo-Peremarch P, Tompalski P, Coops NC, Ruiz LA, 2018. Characterizing understory vegetation in Mediterranean forests using full-waveform airborne laser scanning data. *Remote Sens Environ* 217: 400-413. <https://doi.org/10.1016/j.rse.2018.08.033>

Cubbage F, Harou P, Sills E, 2007. Policy instruments to enhance multi-functional forest management. *Forest Policy Econ* 9: 833-851. <https://doi.org/10.1016/j.forpol.2006.03.010>

Datt B, McVicar TR, Van Niel TG, Jupp DLB, Pearlman JS, 2003. Preprocessing EO-1 Hyperion hyperspectral data to support the application of agricultural indexes. *IEEE T Geosci Remote* 41: 1246-1259. <https://doi.org/10.1109/TGRS.2003.813206>

De Santis A, Chuvieco E, 2007. Burn severity estimation from remotely sensed data: performance of simulation versus empirical models. *Remote Sens Environ* 108: 422-435. <https://doi.org/10.1016/j.rse.2006.11.022>

De Santis A, Chuvieco E, 2009. GeoCBI: A modified version of the Composite Burn Index for the initial assessment of the short-term burn severity from remotely sensed data. *Remote Sens Environ* 113: 554-562. <https://doi.org/10.1016/j.rse.2008.10.011>

Debouk H, Riera-Tatche R, Vega-García C, 2013. Assessing Post-Fire Regeneration in a Mediterranean Mixed Forest Using LiDAR Data and Artificial Neural Networks. 2013. *Photogramm Eng Rem S* 79 (12): 1121-1130. <https://doi.org/10.14358/PERS.79.12.1121>

Díaz-Balteiro L, Romero C, 2008. Making forestry decisions with multiple criteria: A review and an assessment. *Forest Ecol Manag* 255: 3222-3241. <https://doi.org/10.1016/j.foreco.2008.01.038>

Díaz-Delgado R, Pons X, 1999. Empleo de imágenes de teledetección para el análisis de los niveles de severidad causados por el fuego. *Revista de Teledetección* 12: 63-68.

Dobson MC, Ulaby T, Le Toan T, Beaudoin A, Kasischke ES, 1992. Dependence of radar backscatter on coniferous forest biomass. *IEEE Trans Geosci Rem Sens* 30: 412-415. <https://doi.org/10.1109/36.134090>

Domingo D, Lamelas-Gracia MT, Montealegre-Gracia AL, de la Riva-Fernández J, 2017. Comparison of regression models to estimate biomass losses and CO2 emissions using low-density airborne laser scanning data in a burnt Aleppo pine forest. *Eur J Remote Sens* 50 (1): 384-396. <https://doi.org/10.1080/22797254.2017.1336067>

Domingo D, Lamelas M, Montealegre A, de la Riva J, 2018. Estimation of Total Biomass in Aleppo Pine Forest Stands Applying Parametric and Nonparametric Methods to Low-Density Airborne Laser Scanning Data. *Forests* 9 (4): 158. <https://doi.org/10.3390/f9040158>

Duncanson LI, Neumann KO, Wulder MA, 2010. Integration of GLAS and Landsat TM data for aboveground biomass estimation. *Can J Remote Sens* 36 (2): 129-141. <https://doi.org/10.5589/m10-037>

Estornell J, Ruiz LA, Velázquez-Martí B, Fernández-Sarría A, 2011a. Estimation of shrub biomass by airborne LiDAR data in small forest stands. *Forest Ecol Manag* 262: 1697-1703. <https://doi.org/10.1016/j.foreco.2011.07.026>

Estornell J, Ruiz LA, Velázquez-Martí B, 2011b. Study of shrub cover and height using LiDAR data in a Mediterranean area. *For Sci* 57 (3): 171-179.

Estornell J, Ruiz LA, Velázquez-Martí B, Hermsilla T, 2012. Estimation of biomass and volumen of shrub vegetation using LiDAR and spectral data in a Mediterranean environment. *Biomass Bioenerg* 46: 710-721. <https://doi.org/10.1016/j.biombioe.2012.06.023>

EC (European Comission), 2017. https://ec.europa.eu/food/plant/plant_health_biosecurity/legislation/emergency_measures_en

Fassnacht FE, Latifi H, Stereńczak K, Modzelewska A, Lefsky M, Waser LT, Straub C, Ghosh A, 2016. Review of studies on tree species classification from remotely sensed data. *Remote Sens Environ* 186: 64-87. <https://doi.org/10.1016/j.rse.2016.08.013>

Fauzi A, Skidmore AK, Gils H, Schlerf M, Heitkönig I, 2013. Shrimp pond effluent dominates foliar nitrogen in disturbed mangroves as mapped using hyperspectral imagery. *Marine Poll Bull* 76: 42-51. <https://doi.org/10.1016/j.marpolbul.2013.09.033>

Fernández-García V, Santamarta M, Fernández-Manso A, Quintano C, Marcos E, Calvo L, 2018. Burn severity metrics in fire-prone pine ecosystems along a climatic gradient using Landsat imagery, *Remote Sens Environ* 206: 205-217. <https://doi.org/10.1016/j.rse.2017.12.029>

Fernández-Guisuraga JM, Sanz-Ablanedo E, Suárez-Seoane S, Calvo L, 2018. Using Unmanned Aerial Vehicles in Postfire Vegetation Survey Campaigns through Large and Heterogeneous Areas: Opportunities and Challenges. *Sensors* 18: 586. <https://doi.org/10.3390/s18020586>

Fernández-Landa A, Tomé JL, Sandoval VJ, Vallejo R, 2017. Integrando datos LiDAR, información satelital y parcelas del Inventario Forestal Nacional Español en la predicción de variables de inventario. 7º Congreso Forestal Español. 26-30 de junio de 2017. Plasencia, Cáceres, España.

Fernández-Landa A, Fernández-Moya J, Tomé JL, Algeet-Abarquero N, Guillén-Climent ML, Vallejo R, Sandoval V, Marchamalo M, 2018. High resolution forest inventory of pure and mixed stands at regional level combining National Forest Inventory field plots, Landsat, and low density LiDAR. *Int J Remote Sens* 39 (14):4830-4844. <https://doi.org/10.1080/01431161.2018.1430406>

Fernández-Manso O, Fernández-Manso A, Quintano C, 2014. Estimation of aboveground biomass in Mediterranean forests by statistical modelling of ASTER fraction images. *Int J of Appl Earth Obs* 31: 45-56. <https://doi.org/10.1016/j.jag.2014.03.005>

Fernández-Manso A, Quintano C, Roberts DA, 2016a. Burn severity influence on post-fire vegetation cover resilience from Landsat MESMA fraction images time series in Mediterranean forest ecosystems. *Remote Sens Environ* 184: 112-123. <https://doi.org/10.1016/j.rse.2016.06.015>

Fernández-Manso A, Fernández-Manso O, Quintano C, 2016b. SENTINEL-2A red-edge spectral indices suitability for discriminating burn severity. *Int J of Appl Earth Obs* 50: 170-175. <https://doi.org/10.1016/j.jag.2016.03.005>

García M, Riaño D, Chuvieco E, Danson FM, 2010. Estimating biomass carbon stocks for a Mediterranean forest in central Spain using LiDAR height and intensity data. *Remote Sens. Environ.* 114: 816-830. <https://doi.org/10.1016/j.rse.2009.11.021>

García M, Riaño D, Chuvieco E, Salas J, Danson FM, 2011. Multispectral and LiDAR data fusion for fuel type mapping using Support Vector Machine and decision rules. *Remote Sens Environ* 115: 1369-1379. <https://doi.org/10.1016/j.rse.2011.01.017>

García M, Popescu S, Riaño D, Zhao K, Neuenschwander A, Agca M, Chuvieco E, 2012. Characterization of canopy fuels using ICESat/GLAS data, *Remote Sens Environ* 123: 81-89. <https://doi.org/10.1016/j.rse.2012.03.018>

García-Álvarez D, Camacho Olmedo MT, 2017. Changes in the methodology used in the production of the Spanish CORINE: uncertainty analysis of the new maps. *Int J of Appl Earth Obs* 63: 55-67. <https://doi.org/10.1016/j.jag.2017.07.001>

García-Gutiérrez J, González-Ferreiro E, Riquelme-Santos JC, Miranda D, Diéguez-Aranda U, Navarro-Cerrillo RM, 2014. Evolutionary feature selection to estimate forest stand variables using LiDAR, *Int J of Appl Earth Obs* 26: 119-131. <https://doi.org/10.1016/j.jag.2013.06.005>

Garestier F, Dubois-Fernández PC, Papathanassiou KP, 2008. Pine Forest Height Inversion Using Single-Pass X-Band PolInSAR Data. *IEEE Trans Geosci Rem Sens* 46: 59-68. <https://doi.org/10.1109/TGRS.2007.907602>

Gastón A, Ciudad C, Mateo-Sánchez MC, García-Viñas JI, López-Leiva C, Fernández-Landa A, Marchamalo M, Cuevas J, de la Fuente B, Fortin M-J, Saura S, 2017. Species' habitat use inferred from environmental variables at multiple scales: How much we gain from high-resolution vegetation data? *Int J of Appl Earth Obs* 55: 1-8. <https://doi.org/10.1016/j.jag.2016.10.007>

Gómez C, Wulder MA, Montes F, Delgado JA, 2011. Forest structural diversity characterization in Mediterranean pines of Central Spain with QuickBird-2 imagery and canonical correlation analysis. *Can J Remote Sens* 37 (6): 628-642. <https://doi.org/10.5589/m12-005>

Gómez C, Wulder MA, White JC, Montes F, Delgado JA, 2012a. Characterizing 25 years of change in the area, distribution, and carbon stock of Mediterranean pines in Central Spain. *Int J Remote Sens* 33 (17): 5546-5573. <https://doi.org/10.1080/01431161.2012.663115>

Gómez C, Wulder JA, Montes F, Delgado JA, 2012b. Modeling Forest Structural Parameters in the Mediterranean Pines of Central Spain using QuickBird-2 Imagery and Classification and Regression Tree Analysis (CART). *Remote Sens-Basel* 4 (1): 135-159. <https://doi.org/10.3390/rs4010135>

Gómez C, White JC, Wulder MA, Alejandro P, 2014. Historical forest biomass dynamics modelled with Landsat spectral trajectories. *ISPRS J Photogramm93*: 14-28.

Gómez C, Aulló-Maestro I, Montes F, 2016a. Dominant tree species dynamics informed by 30 years of Landsat time series in mountain areas of Northern Spain. *ForestSAT 2016, Santiago (Chile), November*, 15-18.

Gómez C, White JC, Wulder MA, 2016b. Optical remotely sensed time series data for land cover classification: A review. *ISPRS Int J Remote Sens* 116: 55-72. <https://doi.org/10.1016/j.isprsjprs.2016.03.008>

Gómez C, Green D, 2017. Small unmanned airborne systems to support oil and gas pipeline monitoring and mapping. *Arab J Geosci* 10 (9): 202. <https://doi.org/10.1007/s12517-017-2989-x>

Gómez C, Hermosilla T, Martínez-Fernández J, Montes F, Aulló-Maestro I, White JC, Wulder MA, Coops NC, Vázquez A, 2017. Annual cartography of fire (1985-2015) in forest areas of the NW Spain mapped with time series of Landsat data and Composite2Change. *Nuevas plataformas y sensores de teledetección, XVII Congreso de la Asociación Española de Teledetección (Eds. Ruiz LA, Estornell J, Erena M), Murcia (Spain), October 3-7, pp: 169-172.*

Gómez C, Aulló-Maestro I, Alejandro P, Montes F, 2018. Presence of European beech in its Spanish southernmost limit characterized with Landsat intra-annual time series. *AIT2018 IX Conference of the Italian Society of Remote Sensing. Florence (Italy), July 4-6.*

Gonçalves-Seco L, González-Ferreiro E, Diéguez-Aranda U, Fraga-Bugallo B, Crecente R, Miranda D, 2011. Assessomg the attributes of high-density Eucalyptus globulus stands using airborne laser scanner data. *Int J Remote Sens* 32(24): 9821-9841. <https://doi.org/10.1080/01431161.2011.593583>

González-Alonso F, Merino-de-Miguel S, Roldán-Zamarron A, García Gigorro S, Cuevas JM, 2006. Forest biomass estimation through NDVI composites. The role of remote sensed data to assess Spanish forests as carbon sinks. *Int J Remote Sens*, 27: 5409-5415. <https://doi.org/10.1080/01431160600830748>

González-Ferreiro E, Diéguez-Aranda U, Miranda D, 2012. Estimation of stand variables in Pinus radiata D. Don plantations using different LiDAR pulse densities. *Forestry* 85: 281-292. <https://doi.org/10.1093/forestry/cps002>

González-Ferreiro E, Miranda D, Barreiro-Fernández L, Buján S, García-Gutiérrez J, Diéguez-Aranda U, 2013a. Modelling stand biomass fractions in Galician Eucalyptus globulus plantations by use of different LiDAR pulse densities. *Forest Syst* 22: 510-525. <https://doi.org/10.5424/fs/2013223-03878>

González-Ferreiro E, Diéguez-Aranda U, Barreiro-Fernández L, Buján S, Barbosa M, Suárez JC, Bye IJ, Miranda D, 2013b. A mixed pixel- and region-based approach for using airborne laser scanning data for individual tree crown delineation in Pinus radiata D. Don plantations. *Int J Remote Sens* 34(21): 7671-7690. <https://doi.org/10.1080/01431161.2013.823523>

González-Ferreiro E, Diéguez-Aranda U, Crecente-Campo F, Barreiro-Fernández L, Miranda D, Castedo-Dorado F, 2014. Modelling canopy fuel variables for Pinus radiata D. Don in NW Spain with low-density LiDAR data. *I J Wild Fire* 23(3): 350-362. <https://doi.org/10.1071/WF13054>

González-Ferreiro F, Arellano-Pérez S, Castedo-Dorado F, Hevia A, Vega JA, Vega-Nieva D, Álvarez-González JG, Ruiz-González AD, 2017. Modelling the vertical distribution of canopy fuel load using national forest inventory and low-density airborne laser scanning data. *PLoS ONE* 12(4): e0176114. <https://doi.org/10.1371/journal.pone.0176114>

González-Olabarría JR, Rodríguez F, Fernández-Landa A, Mola-Yudego B, 2012. Mapping fire risk in

the model forest of Urbión (Spain) based on airborne LiDAR measurements. For *Ecol Manag* 282: 149-156.

Gonzalo J, López D, Domínguez D, García A, Escapa A, 2017. On the capabilities and limitations of high altitude pseudo-satellites. *Prog Aero Sci* 98: 34-56.

Guerra-Hernández J, Görgens EB, García-Gutiérrez J, Carlos L, Rodríguez E, Tomé M, González-Ferreiro E, 2016. Comparison of ALS based models for estimating aboveground biomass in three types of Mediterranean forest. *Eur. J. Remote Sens* 49: 185-204. <https://doi.org/10.5721/EuJRS20164911>

Guerra-Hernández J, González-Ferreiro E, Monleón VJ, Faias SP, Tomé M, Díaz-Varela RA, 2017. Use of Multi-Temporal UAV-Derived Imagery for Estimating Individual Tree Growth in *Pinus pinea* Stands. *Forests* 8: 300. <https://doi.org/10.3390/f8080300>

Hall RJ, Castilla G, White JC, Cooke BJ, Skakun RS, 2016. Remote sensing of forest pest damage: a review and lessons learned from a Canadian perspective. *Can Entomol* 148: 1-61. <https://doi.org/10.4039/tce.2016.11>

Hajnsek I, Pottier E, Cloude SR, 2003. Inversion of Surface Parameters From Polarimetric SAR. *IEEE Trans Geosci Rem Sens* 41: 727-744. <https://doi.org/10.1109/TGRS.2003.810702>

Henderson FM, Lewis AJ, 1998. Principles and applications of imaging radar. Manual of remote sensing: Third edition, Volume 2. United States. 896 pp.

Hermosilla T, Díaz-Manso JM, Ruiz LA, Recio JA, Fernández-Sarría A, Ferradáns-Nogueira P, 2012. Analysis of parcel-based image classification methods for monitoring the activities of the Land Bank of Galicia (Spain). *Appl Geomat* 4(4): 245-255. <https://doi.org/10.1007/s12518-012-0087-z>

Hermosilla T, Wulder MA, White JC, Coops NC, Hobart GW, 2015. An Integrated Landsat Time Series Protocol for Change Detection and Generation of Annual Gap-Free Surface Reflectance Composites. *Remote Sens Environ* 158: 220-234. <https://doi.org/10.1016/j.rse.2014.11.005>

Hermosilla T, Wulder MA, White JC, Coops NC, Hobart GW, Campbell LB, 2016. Mass data processing of time series Landsat imagery: pixels to data products for forest monitoring. *Int J Digit Earth* 9 (11): 1035-1054. <https://doi.org/10.1080/17538947.2016.1187673>

Hermosilla T, Wulder MA, White JC, Coops NC, Hobart GW, 2017. Updating Landsat time series of surface-reflectance composites and forest change products with new observations. *Int J Appl Earth Obs* 63: 104-111. <https://doi.org/10.1016/j.jag.2017.07.013>

Hermosilla T, Wulder MA, White JC, Coops NC, Hobart GW, 2018. Disturbance-informed annual land cover classification maps of Canada's forested ecosystems for a 29-year Landsat time series. *Can J Remote Sens* 44 (1): 67-87. <https://doi.org/10.1080/07038992.2018.1437719>

Hernández L, Martínez-Fernández J, Cañellas I, Vázquez de la Cueva A, 2014. Assessing spatio-temporal rates, patterns and determinants of biological invasions in forest ecosystems. The case of *Acacia* species in NW Spain, *Forest Ecol Manag* 329: 206-213. <https://doi.org/10.1016/j.foreco.2014.05.058>

Hernández-Clemente R, North PRJ, Hornero A, Zarco-Tejada PJ, 2017. Assessing the effects of forest health on sun-induced chlorophyll fluorescence using the FluorFLIGHT 3-D radiative transfer model

to account for forest structure. *Remote Sens Environ* 193: 165-179. <https://doi.org/10.1016/j.rse.2017.02.012>

Hernando A, Velázquez J, Valbuena R, Legrand M, García-Abril A, 2017. Influence of the Resolution of Forest Cover Maps in Evaluating Fragmentation and Connectivity to Assess Habitat Conservation Status. *Ecol Indic* 79: 295-302. <https://doi.org/10.1016/j.ecolind.2017.04.031>

Hernando A, Puerto L, Mola-Yudego B, Manzanera J, García-Abril A, Maltamo M, Valbuena R, 2019. Estimation of forest biomass components through airborne LiDAR and multispectral sensors. *iForest*

Hevia A, Álvarez-González JG, Ruiz Fernández E, Prendes C, Ruiz González AD, Majada J, González-Ferreiro E, 2016. Modelling canopy fuel and forest stand variables and characterizing the influence of thinning in the stand structure using airborne LiDAR. *Revista Teledetección* 45: 41-55. <https://doi.org/10.4995/raet.2016.3979>

Hilker T, Wulder MA, Coops NC, 2008. Update of forest inventory data with LiDAR and high spatial resolution satellite imagery. *Can J Remote Sens* 34: 5-12. <https://doi.org/10.5589/m08-004>

Huang H, Roy DO, Boschetti L, Zhang HK, Yan L, Kumar SS, Gómez-Dans J, Li J, 2016. Separability analysis of Sentinel-2A multi-spectral instrument (MSI) data for burned area discrimination. *Remote Sens-Basel* 8: 973. <https://doi.org/10.3390/rs8100873>

Huesca M, Merino-de-Miguel S, González-Alonso F, 2013a. An intercomparison of satellite burned area maps derived from MODIS, MERIS, SPOT-VEGETATION and ATSR images. An application to the August 2006 Galicia (Spain) forest fires. *For Syst* 22 (2): 222-231.

Huesca M, Merino-de-Miguel S, González-Alonso F, Martínez S, Cuevas JM, Calle A, 2013b. Using AHS hyper-spectral images to study forest vegetation recovery after a fire. *Int J Remote Sens* 34 (11): 4025-4048. <https://doi.org/10.1080/01431161.2013.772313>

Hyypä J, Inkinen M, 1999. Detecting and estimating attributes for single tree using laser scanner. *Phot J Fin* 16: 27-42.

INE 2017. España en cifras. https://www.ine.es/prodyser/espa_cifras/2017/index.html#5

Jensen J, 2005. *Introductory digital image processing: a remote sensing perspective*, 3rd ed. Pearson Education, Inc. 526 pp.

Joshi N, Mitchard ETA, Brolly M, Schumacher J, Fernández-Landa A, Johannsen VK, Marchamalo M, Fensholt R, 2017. Understanding 'saturation' of radar signals over forests. *Sci Rep-UK* 7 (1): 3505. <https://doi.org/10.1038/s41598-017-03469-3>

Karnkowski W, Sahajdak A, 2010. Occurrence of the pinewood nematode in Portugal and Spain - Threat for pine forests in Europe. *Prog Plant Prot* 50: 1260-1264.

Keane RE, Burgan R, Wagtendonk JV, 2001. Mapping wildland fuels for fire management across multiple scales: Integrating remote sensing, GIS, and biophysical modeling, *Int J Wildland Fire* 10: 301-319. <https://doi.org/10.1071/WF01028>

Kennedy RE, Yang Z, Cohen WB, 2010. Detecting trends in forest disturbance and recovery using yearly Landsat time series: 1. LandTrendr-temporal segmentation algorithms. *Remote Sens Environ* 114: 2897-2910. <https://doi.org/10.1016/j.rse.2010.07.008>

Kennedy RE, Yang Z, Braaten J, Copass C, Antonova N, Jordan C, Nelson P, 2015. Attribution of disturbance change agent from Landsat time-series in support of habitat monitoring in the Puget Sound region, USA. *Remote Sens Environ* 166: 271-285. <https://doi.org/10.1016/j.rse.2015.05.005>

Key CH, Benson NC, 1999. Measuring and remote sensing of burn severity. In L. F. Neuenschwander and K. C. Ryan (Eds.), *Proceedings Joint Fire Science Conference and Workshop, Vol. II.* (pp. 284). Moscow, ID: University of Idaho and International Association of Wildland Fire.

Kissinger G, Herold M, De Sy V, 2012. *Drivers of Deforestation and Forest Degradation: A Synthesis Report for REDD+ Policymakers.* Lexeme Consulting, Vancouver Canada, 46 pp.

Le Toan T, Beaudoin A, Guyon D, 1992. Relating forest biomass to SAR data. *IEEE Trans Geosci Rem Sens* 30: 403-411. <https://doi.org/10.1109/36.134089>

Le Toan T, Quegan S, Davidson MWJ, Balzter H, Paillou P, Papathanassiou K, Plummer S, Rocca F, Saatchi S, Shugart H, Ulander LMH, 2011. The BIOMASS mission: Mapping global forest biomass to better understand the terrestrial carbon cycle. *Remote Sens Environ* 115: 2850-2860. <https://doi.org/10.1016/j.rse.2011.03.020>

Leberl F, Irschara A, Pock T, Meixner P, Gruber M, Scholz S, Wiechert A, 2010. Point clouds: LiDAR versus three dimensional vision. *Photogramm Eng Rem Sens* 76: 1123-1134.

Lefsky MA, Cohen WB, Acker SA, Parker GG, Spies TA, Harding D, 1999. LiDAR remote sensing of the canopy structure and biophysical properties of Douglas-fir western hemlock forests. *Remote Sens Environ* 70: 339-361. [https://doi.org/10.1016/S0034-4257\(99\)00052-8](https://doi.org/10.1016/S0034-4257(99)00052-8)

Lefsky MA, 2010. A global forest canopy height map from the Moderate Resolution Imaging Spectroradiometer and the Geoscience Laser Altimeter System. *Geophys Res Lett* 37 L15401. <https://doi.org/10.1029/2010GL043622>

Li J, Roy DP, 2017. A global analysis of Sentinel-2A, Sentinel-2B and Landsat-8 data revisit intervals and implications for terrestrial monitoring. *Remote Sens-Basel* 9: 902. <https://doi.org/10.3390/rs9090902>

Lisein J, Pierrot-Deseilligny M, Bonnet, S, Lejeune P, 2013. A Photogrammetric Workflow for the Creation of a Forest Canopy Height Model from Small Unmanned Aerial System Imagery. *Forests* 4: 922-944. <https://doi.org/10.3390/f4040922>

Lohberger S, Stängel M, Atwood EC, Siegert F, 2018. Spatial evaluation of Indonesia's 2015 fire-affected area and estimated carbon emissions using Sentinel-1. *Glob Chang Biol* 24 (2): 644-654. <https://doi.org/10.1111/gcb.13841>

López-Sánchez JM, Ballester-Berman JD, 2009. Potentials of Polarimetric SAR Interferometry for Agriculture Monitoring. *Radio Sci* 44: RS2010. <https://doi.org/10.1029/2008RS004078>

Malak DA, Pausas JG, Pardo-Pascual JE, Ruiz LA, 2015. Fire recurrence and the dynamics of the Enhanced Vegetation Index in a Mediterranean ecosystem. *Int J App Geosp Res* 6 (2): 18-35. <https://doi.org/10.4018/ijagr.2015040102>

Manfreda S, McCabe MF, Miller PE, Lucas R, Pajuelo Madrigal V, Mallinis G, Ben Dor E, Helman D,

Estes L, Ciraolo G, et al., 2018. On the Use of Unmanned Aerial Systems for Environmental Monitoring. *Remote Sens-Basel* 10: 641. <https://doi.org/10.3390/rs10040641>

Manzanera JA, García-Abril A, Pascual C, Tejera R, Martín-Fernández S, Tokola T, Valbuena R, 2016. Fusion of airborne LiDAR and multispectral sensors reveals synergic capabilities in forest structure characterization. *GISci Rem Sens* 53(6): 723-738. <https://doi.org/10.1080/15481603.2016.1231605>

MAPAMA, 2011. Anuario de estadística forestal. Ministerio de Agricultura, Alimentación y Medio Ambiente, Spain, 103 pp

Marino E, Ranz P, Tomé JL, Noriega MA, Esteban J, Madrigal J, 2016. Generation of high-resolution fuel models from discrete airborne laser scanner and Landsat-8 OLI: a low-cost and highly updated methodology for large areas. *Remote Sens Environ* 187: 267-280. <https://doi.org/10.1016/j.rse.2016.10.020>

Marino E, Tomé JL, Madrigal J, Guijarro M, Hernando C, 2017a. Efecto de la densidad de pulsos LiDAR en la caracterización estructural de combustibles en masas de pinar. 7º Congreso Forestal Español. 26-30 de junio de 2017. Plasencia, Cáceres, España.

Marino E, Ranz P, Tomé JL, 2017b. Evolución post-incendio de la estructura de la vegetación en el PN de Garajonay a partir de datos LiDAR. 7º Congreso Forestal Español. 26-30 de junio de 2017. Plasencia, Cáceres, España.

Marino E, Montes F, Tomé JL, Navarro JA, Hernando C, 2018. Vertical forest structure analysis for wildfire prevention: comparing airborne laser scanning data and stereoscopic hemispherical images. *Int J Appl Earth* 73: 438-449. <https://doi.org/10.1016/j.jag.2018.07.015>

Martín-Alcón S, Coll LI, de Cáceres M, Guitart L, Cabré M, Just A, González-Olabarría JR, 2015. Combining aerial LiDAR and multi-spectral imagery to assess post-fire regeneration types in a Mediterranean forest. *Can J For Res* 45 (7): 856-866. <https://doi.org/10.1139/cjfr-2014-0430>

Martínez S, Chuvieco E, Aguado I, Salas J, 2017. Severidad y regeneración en grandes incendios forestales: análisis a partir de series temporales de imágenes Landsat. *Revista de Teledetección* 49: 17-32. <https://doi.org/10.4995/raet.2017.7182>

Martínez-Fernández J, Ruiz-Benito P, Bonet-Jornet A, Gómez C, 2019. Methodological variations in the production of CORINE Land Cover and consequences for long-term land cover change studies. The case of Spain. *Int J Remote Sens*

Matasci G, Hermosilla T, Wulder MA, White JC, Coops NC, Hobart GW, Zald HSJ, 2018. Large-area mapping of Canadian boreal forest cover, height, biomass and other structural attributes using Landsat composites and LiDAR plots. *Remote Sens Environ* 209: 90-106. <https://doi.org/10.1016/j.rse.2017.12.020>

Mauro F, Valbuena R, Manzanera JA, García-Abril A, 2011. Influence of Global Navigation Satellite System errors in positioning inventory plots for tree-height distribution studies. *Can J For Res* 41 (1): 11-23. <https://doi.org/10.1139/X10-164>

Mauro F, Molina I, García-Abril A, Valbuena R, Ayuga-Téllez E, 2016. Remote sensing estimates and measures of uncertainty for forest variables at different aggregation levels. *Environmetrics* 27 (4): 225-238. <https://doi.org/10.1002/env.2387>

Mauro F, Monleón VJ, Temesgen H, Ford KR, 2017a. Analysis of area level and unit level models for small area estimation in forest inventories assisted with LiDAR auxiliary information. *PloS ONE* 12 (12), e0189401 <https://doi.org/10.1371/journal.pone.0189401>

Mauro F, Monleón VJ, Temesgen H, Ruiz LA, 2017b. Analysis of spatial correlation in predictive models of forest variables that use LiDAR auxiliary information. *Can J For Res* 47 (6): 788-799. <https://doi.org/10.1139/cjfr-2016-0296>

Melicharová L, Vizoso-Arribe O, 2012. Situation of sweet chestnut (*Castanea sativa* Mill.) in Spain, Galicia: A review. *Scientia Agr Boh* 2012: 78-84.

Méndez E, Valés JJ, Pino I, Granado L, Montoya G, Prieto R, Carpintero IR, Giménez de Azcárate F, Cáceres F, Moreira JM, et al., 2016. Determination of forest biomass using remote sensing techniques with radar images. Pilot study in area of the province of Huelva. *REDIAM. Revista de Teledetección* 45: 71-86. <https://doi.org/10.4995/raet.2016.3984>

Merino de Miguel S, Huesca M, González-Alonso F, 2010. MODIS reflectance and active fire data for burn mapping and assessment at regional level. *Ecol Model* 221: 67-74. <https://doi.org/10.1016/j.ecolmodel.2009.09.015>

Moessner KE, 1953. Photo interpretation in forest inventories. *Photogramm Eng*, June 1953: 496-507.

Montealegre AL, Lamelas MT, de la Riva J, García-Martín A, Escribano F, 2016. Use of low point density ALS data to estimate stand-level structural variables in Mediterranean Aleppo pine forest. *Forestry* 0: 1-10. <https://doi.org/10.1093/forestry/cpw008>

Montealegre AL, Lamelas MT, Tanase MA, de la Riva J, 2017a. Estimación de la severidad en incendios forestales a partir de datos LIDAR-PNOA y valores de Composite Burn Index. *Revista de Teledetección* 49: 1-16. <https://doi.org/10.4995/raet.2017.7371>

Montealegre AL, Lamelas-Gracia MT, García-Martín A, de la Riva-Fernández J, Escribano-Bernal F, 2017b. Using low density discrete Airborne Laser Scanning data to assess the potential carbon dioxide emission in case of a fire event in a Mediterranean pine forest. *GISci Rem Sens* 54 (5): 721-740. <https://doi.org/10.1080/15481603.2017.1320863>

Montero G, Ruiz-Peinado R, Muñoz M, 2005. Producción de biomasa y fijación de CO2 por los bosques espa-oles. Monografías Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria, Serie Forestal, Madrid, Spain.

Montero G, Serrada R, 2013. La situación de los bosques y el sector forestal en España-ISFE 2013. Sociedad Española de Ciencias Forestales. Lourizán (Pontevedra), Spain. 257 pp.

Moreira A, Krieger G, Hajnsek I, Papathanassiou K, Younis M, Lopez-Dekker P, Huber S, Villano M, Pardini M, Eineder M, et al., 2015. Tandem-L/ALOSNext: A Highly Innovative Bistatic SAR Mission for Global Observation of Dynamic Processes on the Earth's Surface. *IEEE Geosc Remote Sens Mag* 3 (2): 8-23. <https://doi.org/10.1109/MGRS.2015.2437353>

Næsset E, 2002. Predicting forest stand characteristics with airborne scanning laser using a practical two-stage procedure and field data. *Remote Sens Environ* 80: 88-99. [https://doi.org/10.1016/S0034-4257\(01\)00290-5](https://doi.org/10.1016/S0034-4257(01)00290-5)

Navarro JA, Fernández-Landa A, Tomé JL, Guillén-Climent ML, Ojeda JC, 2018. Testing the quality of forest variable estimation using dense image matching: a comparison with airborne laser scanning in a Mediterranean pine forest. *Int J Rem Sens* 39 (14): 4744-4760. <https://doi.org/10.1080/01431161.2018.1471551>

Navarro-Cerrillo RM, González-Ferreiro E, García-Gutiérrez J, Ceacero Ruiz CJ, Hernández-Clemente R, 2017. Impact of plot size and model selection on forest biomass estimation using airborne LiDAR: A case study of pine plantations in southern Spain. *J For Sci* 63: 88-97. <https://doi.org/10.17221/86/2016-JFS>

Oeser J, Pflugmacher D, Senf C, Heurich M, Hostert P, 2017. Using intra-annual Landsat time series for attributing forest disturbance agents in Central Europe. *Forests* 8: 251. <https://doi.org/10.3390/f8070251>

Olesk A, Praks J, Antropov O, Zalite K, Arumäe T, Voormansik K, 2016. Interferometric SAR coherence models for characterization of hemiboreal forests using TanDEM-X data. *Remote Sens-Basel* 8: 700. <https://doi.org/10.3390/rs8090700>

Packalén P, Maltamo M, 2006. Predicting the plot volume by tree species using airborne laser scanning and aerial photographs. *Forest Sci* 52: 611–622.

Packalén P, Suvanto A, Maltamo M, 2009. A two stage method to estimate species-specific growing stock. *Photogramm Eng Rem Sens* 75: 1451–1460. <https://doi.org/10.14358/PERS.75.12.1451>

Pajares G, 2015. Overview and current status of remote sensing applications based on Unmanned Aerial Vehicles (UAVs). *Photogramm Eng Rem Sens* 81 (4): 281-329. <https://doi.org/10.14358/PERS.81.4.281>

Parra A, Chuvieco E, 2005. Assessing burn severity using Hyperion data. In J Riva, Pérez-Cabello F, Chuvieco E (Eds.) *Proceedings of the 5th international workshop on remote sensing and GIS applications to forest fire management: fire effects assessment* (pp 239-244) Paris. Universidad d Zaragoza, GOFC-GOLD, EARSeL.

Pascual A, Pukkala T, Rodríguez F, de-Miguel S, 2016. Using Spatial Optimization to Create Dynamic Harvest Blocks from LiDAR-Based Small Interpretation Units. *Forests* 7(10): 220. <https://doi.org/10.3390/f7100220>

Pascual A, Pukkala T, de-Miguel S, 2018a. Effects of plot positioning errors on the optimality of harvest prescriptions in spatial forest planning based on ALS data. *Forests* 9(7): 371. <https://doi.org/10.3390/f9070371>

Pascual A, Pukkala T, de-Miguel S, Pesonen A, Packalén P, 2018b. Influence of timber harvesting costs on the layout of cuttings and economic return in forest planning based on dynamic treatment units. *For Syst* 27:1.

Pascual C, García-Abril A, García-Montero LG, Martín-Fernández S, Cohen WB, 2008. Object-based semi-automatic approach for forest structure characterization using LIDAR data in heterogeneous *Pinus sylvestris* stands. *Forest Ecol Manag* 255: 3677-3685. <https://doi.org/10.1016/j.foreco.2008.02.055>

Pascual C, García-Abril A, Cohen WB, Martín-Fernández S, 2010. Relationship between LiDAR-

derived forest canopy height and Landsat images. *Int J Remote Sens* 31 (5): 1261-1280. <https://doi.org/10.1080/01431160903380656>

Pascual C, García-Montero LG, Arroyo LA, García-Abril A, 2013. Increasing the use of expert opinion in forest characterisation approaches based on LiDAR data. *Annals of Forest Science* 70: 87-99. <https://doi.org/10.1007/s13595-012-0232-1>

Pasquarella VJ, Holden CE, Kaufman L, Woodcock CE, 2016. From imagery to ecology: leveraging time series of all available Landsat observations to map and monitor ecosystem state and dynamics. *Rem Sens Ecol Conserv* 2 (3): 151-170. <https://doi.org/10.1002/rse2.24>

Pasquarella VJ, Bradley BA, Woodcock CE, 2017. Near-Real-Time Monitoring of Insect Defoliation Using Landsat Time Series. *Forests* 8 (8): 275. <https://doi.org/10.3390/f8080275>

Pulliainen JT, Heiska K, Hyyappa J, Hallikainen MT, 1994. Backscattering properties of boreal forests at the C- and X-Bands. *IEEE Trans Geosci Rem Sens*, 32, 1041-1050. <https://doi.org/10.1109/36.312892>

Qi W, Dubayah RO, 2016. Combining Tandem-X InSAR and simulated GEDI LiDAR observations for forest structure mapping. *Remote Sens Environ* 187: 253-266. <https://doi.org/10.1016/j.rse.2016.10.018>

Quintano C, Fernández-Manso A, Fernández-Manos O, Shimabukuro YE, 2006. Mapping burned areas in Mediterranean countries using spectral mixture analysis from a uni-temporal perspective. *Int J Remote Sens* 27(4): 645-662. <https://doi.org/10.1080/01431160500212195>

Quintano C, Fernández-Manso A, Calvo L, Marcos E, Valbuena L, 2015. Land Surface temperature as potential indicator of burn severity in forest Mediterranean ecosystems. *Int J Appl Earth* 36: 1-12. <https://doi.org/10.1016/j.jag.2014.10.015>

Quintano C, Fernández-Manso A, Fernández-Manso O, 2018. Combination of Landsat and Sentinel-2 MSI data for initial assessing of burn severity. *Int J Appl Earth Obs Geoinformation* 64: 221-225. <https://doi.org/10.1016/j.jag.2017.09.014>

Radeloff VC, Mildenoff DJ, Boyce MS, 1999. Detecting Jack Pine budworm defoliation using spectral mixture analysis: separating effects from determinants. *Remote Sens Environ* 69: 156-169. [https://doi.org/10.1016/S0034-4257\(99\)00008-5](https://doi.org/10.1016/S0034-4257(99)00008-5)

Regos A, Ninyerola M, Moré G, Pons X, 2015. Linking land cover dynamics with driving forces in mountain landscape of the Northwestern Iberian Peninsula. *Int J Appl Earth* 38: 1-14. <https://doi.org/10.1016/j.jag.2014.11.010>

Riaño D, Chuvieco E, Salas J, Palacios-Orueta A, Bastarrika A, 2002. Generation of fuel type maps from Landsat TM images and ancillary data in Mediterranean ecosystems. *Can J Forest Res* 32: 1301-1315. <https://doi.org/10.1139/x02-052>

Riaño D, Chuvieco E, Condés S, González-Matesanz J, Ustin SL, 2004. Generation of crown bulk density for *Pinus sylvestris* L. from LiDAR. *Remote Sens Environ* 92: 345-352. <https://doi.org/10.1016/j.rse.2003.12.014>

Rignot EJ, Way J, Williams C, Viereck L, 1994. Radar Estimates of Aboveground Biomass in Boreal Forests of Interior Alaska. *IEEE Trans Geosci Rem Sens* 32: 1117-

1124. <https://doi.org/10.1109/36.312903>

Robles A, Rodríguez-Garrido MA, Álvarez-Taboada MF, 2016. Characterization of wildland-urban interfaces using LiDAR data to estimate the risk of wildfire damage. *Revista de Teledetección. (Special Issue)*: 57-69.

Rock BN, Vogelmann JE, Williams DL, Vogehmann AF, Hoshizaki T, 1986. Remote detection of forest damage. *Bioscience* 36: 439-445. <https://doi.org/10.2307/1310339>

Rubio A, Gavilán RG, Montes F, Gutiérrez-Girón A, Díaz-Pines E, Mezquida ET, 2011. Biodiversity measures applied to stand-level management: Can they really be useful? *Ecol Indic* 11: 545-556. <https://doi.org/10.1016/j.ecolind.2010.07.011>

Ruiz LA, Recio JA, Fernández-Sarría A, 2005. Clasificación de entornos forestales mediterráneos mediante técnicas de análisis de texturas. *Cuadernos de la Sociedad Española de Ciencias Forestales (SECF)* 19: 187-192.

Ruiz LA, Hermosilla T, Mauro F, Godino M, 2014. Analysis of the influence of plot size and LiDAR density on forest structure attribute estimates. *Forests* 5 (5): 936-957. <https://doi.org/10.3390/f5050936>

Ruiz LA, Recio JA, Crespo-Peremarch P, Sapena M, 2018. An object-based approach for mapping forest structural types based on low density LiDAR and multispectral imagery. *Geocarto Int* 33: 443-457. <https://doi.org/10.1080/10106049.2016.1265595>

Ruiz-Gallardo JR, Castaño S, Calera A, 2004. Application of remote sensing and GIS to locate priority intervention areas after wildland fires in Mediterranean systems: a case study from south-eastern Spain. *Int J Wildland Fire* 13: 241-252. <https://doi.org/10.1071/WF02057>

Ruiz-Peinado R, Del Rio M, Montero G, 2011. New models for estimating the carbon sink capacity of Spanish softwood species. *For Syst* 20: 176-188.

Rullán-Silva CD, Olthoff AE, Delgado JA, Pajares-Alonso JA, 2013. Remote monitoring of forest insect defoliation. A review. *Forest Syst* 22 (3): 377-391. <https://doi.org/10.5424/fs/2013223-04417>

Rullán-Silva C, Olthoff AE, Pando V, Pajares JA, Delgado JA, 2015. Remote monitoring of defoliation by the beech leaf-mining weevil *Rhynchaenus fagi* in northern Spain. *Forest Ecol Manag* 347: 200-208. <https://doi.org/10.1016/j.foreco.2015.03.005>

Sandberg G, Ulander LMH, Fransson JES, Holmgren J, Toan TL, 2011. L- and P-band backscatter intensity for biomass retrieval in hemiboreal forest. *Remote Sens Environ* 115: 2874-2886. <https://doi.org/10.1016/j.rse.2010.03.018>

Sangüesa-Barreda G, Camarero JJ, García-Martín A, Rodolfo Hernández R, de la Riva J, 2014. Remote-sensing and tree-ring based characterization of forest defoliation and growth loss due to the Mediterranean pine processionary moth. *Forest Ecol Manag* 320: 171-181. <https://doi.org/10.1016/j.foreco.2014.03.008>

Sangüesa-Barreda G, Camarero JJ, Oliva J, Montes F, Gazol A, 2015. Past logging, drought and pathogens interact and contribute to forest dieback. *Agric For Meteorol* 208: 85-94. <https://doi.org/10.1016/j.agrformet.2015.04.011>

Sankey T, Donager J, McVay J, Sankey JB, 2017. UAV LiDAR and hyperspectral fusion for forest monitoring in the southwestern USA. *Remote Sens Environ* 195: 30-43. <https://doi.org/10.1016/j.rse.2017.04.007>

Schlerf M, Atzberger C, Hill J, Buddenbaum H, Werner W, Schuler G, 2010. Retrieval of chlorophyll and nitrogen in Norway spruce (*Picea abies* L. Karst.) using imaging spectroscopy. *Int J Appl Earth 12*: 17-26. <https://doi.org/10.1016/j.jag.2009.08.006>

Schutz BE, Zwally HJ, Shuman CA, Hancock D, DiMarzio JP, 2005. Overview of the ICESat Mission, *Geophys Res Lett* 32: L21S01. <https://doi.org/10.1029/2005GL024009>

Sevillano-Marco E, Fernández-Manso A, Quintano C, Poulain M, 2013. CCD CBERS and ASTER data in dasometric characterization of *Pinus radiata* D. Don (North-Western Spain). *Cerne, Lavras* 19(1): 103-110.

Shimada M, Itoh T, Motooka T, Watanabe M, Shiraishi T, Thapa R, Lucas R, 2014. New global forest/non-forest maps from ALOS PALSAR data (2007–2010). *Remote Sens Environ* 155: 13-31. <https://doi.org/10.1016/j.rse.2014.04.014>

Sherald J, 2007. Bacterial Leaf Scorch of Landscape Trees: What We Know and What We Do Not Know. *Arb Urb Forestry* 33.

Simonson W, Allen H, Coomes D, 2018. Effect of Tree Phenology on LiDAR Measurement of Mediterranean Forest Structure. *Remote Sens* 10: 659 <https://doi.org/10.3390/rs10050659>

Silveira EM, de Mello JM, Acerbi FW, dos Reis AA, Withey KD, Ruiz LA, 2018. Characterizing landscape spatial heterogeneity using semivariogram parameters derived from NDVI images. *CERNE*, 23 (4): 413-422. <https://doi.org/10.1590/01047760201723042370>

Smith MW, Carrivick J, Quincey D, 2016. Structure from Motion Photogrammetry in Physical Geography. *Prog Phys Geog* 40 (2): 247-275. <https://doi.org/10.1177/0309133315615805>

Tanase M, Santoro M, Wegmüller U, de la Riva J, Pérez-Cabello F, 2010b. Properties of X-, C- and L-band repeat-pass interferometric SAR coherence in Mediterranean pine forests affected by fires, *Remote Sens Environ* 114: 2182-2194. <https://doi.org/10.1016/j.rse.2010.04.021>

Tanase MA, de la Riva J, Pérez-Cabello F, 2011a. Estimating burn severity at the regional level using optically based indices. *Can J For Res* 41: 863-872. <https://doi.org/10.1139/x11-011>

Tanase MA, de la Riva J, Santoro M, Pérez-Cabello F, Kasischke E, 2011b. Sensitivity of SAR data to post-fire forest regrowth in Mediterranean and boreal forests. *Remote Sens Environ* 115: 2075-2085. <https://doi.org/10.1016/j.rse.2011.04.009>

Tanase MA, Panciera R, Lowell K, Tian S, García-Martín A, Walker JP, 2014a. Sensitivity of L-band radar backscatter to forest biomass in semi-arid environments: a comparative analysis of parametric and non-parametric models. *IEEE Trans Geosci Rem Sens*: 52, 1-15. <https://doi.org/10.1109/TGRS.2013.2283521>

Tanase MA, Panciera R, Lowell K, Aponte C, Hacker JM, Walker JP, 2014b. Forest biomass estimation at high spatial resolution: Radar vs. LiDAR sensors. *IEEE Trans Geosci Rem Sens Lett* 11 (3): 711-715. <https://doi.org/10.1109/LGRS.2013.2276947>

Tanase MA, Santoro M, Aponte C, De la Riva J, 2014c. Polarimetric Properties of Burned Forest Areas at C- and L-Band. *IEEE Trans Geosci Rem Sens* 7 (1): 267-276.

Tanase MA, Kennedy R, Aponte C, 2015a. Fire severity from space: a comparison of active and passive sensors and their synergy for different forest types. *Int J Wildland Fire* 24(8): 1062-1075. <https://doi.org/10.1071/WF15059>

Tanase MA, Kennedy R, Aponte C, 2015b. Radar Burn Ratio for fire severity estimation at canopy level: an example for temperate forests. *Remote Sens Environ* 170: 14-31. <https://doi.org/10.1016/j.rse.2015.08.025>

Tanase MA, Panciera R, Lowell K, Aponte C, 2015c. Monitoring live fuel moisture in semi-arid environments using L-band radar data. *Int J Wildland Fire* 24: 560-572. <https://doi.org/10.1071/WF14149>

Tanase MA, Aponte C, Mermoz S, Bouvet A, Le Toan T, Heurich M, 2018. Detection of windthrows and insect outbreaks by L-band SAR: A case study in the Bavarian Forest National Park. *Remote Sens Environ* 209: 700-711. <https://doi.org/10.1016/j.rse.2018.03.009>

Tebaldini S, Rocca F, 2012. Multibaseline Polarimetric SAR Tomography of a Boreal Forest at P- and L-Bands. *IEEE T Geosci Remote* 50: 232-246. <https://doi.org/10.1109/TGRS.2011.2159614>

Tomé JL, Esteban J, Martín-Alcón S, Escamochero I, Fernández-Landa A, 2017. Forestmap, inventario forestal online a partir de datos LiDAR en la Región de Murcia. Nuevas plataformas y sensores de teledetección, XVII Congreso de la Asociación Española de Teledetección (Eds. Ruiz LA, Estornell J, Erena M), Murcia (Spain), October 3-7, pp: 147-150.

Tompalski P, Coops NC, Marshall PL, White JC, Wulder MA, Bailey T, 2018. Combining multi-date airborne laser scanning and digital aerial photogrammetric data for forest growth and yield modelling. *Remote Sens-Basel* 10: 347. <https://doi.org/10.3390/rs10020347>

Tomppo E, Olsson H, Stahl G, Nilsson M, Hagner O, Katila M, 2008. Combining national forest inventory field plots and remote sensing data for forest databases. *Remote Sens Environ* 112: 1982-1999. <https://doi.org/10.1016/j.rse.2007.03.032>

Torresan C, Beaton A, Carotenuto F, Filippo S, Gioli B, Matese A, Miglietta F, Magnoli C, Zaldea A, Wallace L, 2017. Forestry applications of UAVs in Europe: a review. *Int J Remote Sens* 38 (8-10): 2427-2447. <https://doi.org/10.1080/01431161.2016.1252477>

Transon J, Andrimont R, Maignard A, Defourny P, 2018. Survey of hyperspectral Earth Observation applications from space in the Sentinel-2 context. *Remote Sens-Basel* 10: 157. <https://doi.org/10.3390/rs10020157>

Trassierra A, Esteban J, Fernández-Landa A, Sabín P, Sánchez-Pellicer T, Tomé JL, 2017. Modelos de estimación de carga de biomasa aérea de matorral a partir de diferentes fuentes de información: LiDAR y Landsat. 7º Congreso Forestal Español. 26-30 de junio de 2017. Plasencia, Cáceres, España.

Turner DP, Cohen WB, Kennedy RE, Fassnacht KS, Briggs JM, 1999. Relationship between leaf area index and Landsat TM spectral vegetation indices across three temperate zone sites. *Remote Sens Environ* 70: 52-68. [https://doi.org/10.1016/S0034-4257\(99\)00057-7](https://doi.org/10.1016/S0034-4257(99)00057-7)

UNECE (United Nations Economic Commission For Europe) ICP Forests Programme Co-ordinating

Centre (Ed.), 2016, Manual on Methods and Criteria for Harmonized Sampling, Assessment, Monitoring and Analysis of the Effects of Air Pollution on Forests. Thunen Institute of Forest Ecosystems, Eberswalde. <http://icp-forests.net/page/icp-forests-manual>

Ungar S, Pearlman J, Mendenhall J, Reuter D, 2003. Overview of the Earth Observing-1 (EO-1) mission. *IEEE T Geosci Remote* 41: 1149–1159. <https://doi.org/10.1109/TGRS.2003.815999>

Valbuena R, Mauro F, Arjonilla FJ, Manzanera JA, 2011. Comparing Airborne Laser Scanning-Imagery Fusion Methods Based on Geometric Accuracy in Forested Areas. *Remote Sens Environ* 115(8): 1942-1956. <https://doi.org/10.1016/j.rse.2011.03.017>

Valbuena R, Mauro F, Rodríguez-Solano R, Manzanera JA, 2012. Partial Least Squares for Discriminating Variance Components in GNSS Accuracy Obtained Under Scots Pine Canopies. *Forest Sci* 58(2): 139-153. <https://doi.org/10.5849/forsci.10-025>

Valbuena R, De Blas A, Martín Fernández S, Maltamo M, Nabuurs GJ, Manzanera JA, 2013a. Within-Species Benefits of Back-projecting Laser Scanner and Multispectral Sensors in Monospecific *P. sylvestris* Forests. *Eur J Remote Sens* 46: 401-416. <https://doi.org/10.5721/EuJRS20134629>

Valbuena R, Maltamo M, Martín-Fernández S, Packalen P, Pascual C, Nabuurs G-J, 2013b. Patterns of covariance between airborne laser scanning metrics and Lorenz curve descriptors of tree size inequality. *Can J Remote Sens* 39(1): 18-31. <https://doi.org/10.5589/m13-012>

Valbuena R, Packalen P, García-Abril A, Mehtätalo L, Maltamo M, 2013c. Characterizing Forest Structural Types and Shelterwood Dynamics from Lorenz-based Indicators Predicted by Airborne Laser Scanning. *Can J For Res* 43: 1063-1074. <https://doi.org/10.1139/cjfr-2013-0147>

Valbuena R, Maltamo M, Packalen P, 2016a. Classification of Multi-Layered Forest Development Classes from Low-Density National Airborne LiDAR Datasets. *Forestry* 89: 392-341. <https://doi.org/10.1093/forestry/cpw010>

Valbuena R, Maltamo M, Packalen P, 2016b. Classification of Forest Development Stages from National Low-Density LiDAR Datasets: a Comparison of Machine Learning Methods. *Revista de Teledetección* 45: 15-25. <https://doi.org/10.4995/raet.2016.4029>

Valbuena R, Hernando A, Manzanera JA, Martínez-Falero E, García-Abril A, Mola-Yudego B, 2017a. Most Similar Neighbour Imputation of Forest Attributes Using Metrics Derived from Combined Airborne LiDAR and Multispectral Sensors. *Int J Digit Earth* 11 (12): 1205-1218. <https://doi.org/10.1080/17538947.2017.1387183>

Valbuena R, Hernando A, Manzanera JA, Görgens EB, Almeida DRA, Mauro F, García-Abril A, Coomes DA, 2017b. Enhancing of accuracy assessment for forest above-ground biomass estimates obtained from remote sensing via hypothesis testing and overfitting evaluation. *Eco Mod* 622: 15-26. <https://doi.org/10.1016/j.ecolmodel.2017.10.009>

Valbuena-Rabadán M, Santamaría-Peña J, Sanz-Adán F, 2016. Estimation of diameter and height of individual trees for *Pinus sylvestris* L. based on the individualising of crowns using airborne LiDAR and the National Forest Inventory data. *For Sys* 25(1): e046

Varo-Martínez MA, Navarro-Cerrillo RM, Hernández-Clemente R, Duque-Lazo J, 2017. Semi-automated stand delineation in Mediterranean *Pinus sylvestris* plantations through segmentation of LiDAR data: The influence of pulse density. *Int J Appl Earth Obs* 56: 54-

64. <https://doi.org/10.1016/j.jag.2016.12.002>

Vázquez de la Cueva A, 2008. Structural attributes of three forest types in central Spain and Landsat ETM+ information evaluated with redundancy analysis. *Int J Remote Sens* 29: 5657-5676. <https://doi.org/10.1080/01431160801891853>

Verdú F, Salas J, 2010. Cartografía de áreas quemadas mediante análisis visual de imágenes de satélite en la España peninsular para el periodo 1991–2005. *Geofocus* 10: 54–81.

Viana-Soto A, Aguado I, Martínez S, 2017. Assessment of post-fire vegetation recovery using fire severity and geographical data in the Mediterranean region (Spain). *Environments* 4: 90. <https://doi.org/10.3390/environments4040090>

Vicente-Serrano SG, Pérez-Cabello F, Lasanta T, 2011. *Pinus halepensis* regeneration after a wildfire in a semiarid environment: assessment using multitemporal Landsat images. *Int J Wildland Fire* 20Ñ 195-208.

Viedma O, Quesada J, Torres I, De Santis A, Moreno JM, 2015. Fire severity in a large fire in a *Pinus pinaster* forest is highly predictable from burning conditions, stand structure, and topography. *Ecosystems* 18: 237-250. <https://doi.org/10.1007/s10021-014-9824-y>

Yebra M, Chuvieco E, 2009. Generation of a species-specific look-up table for fuel moisture content assessment. *IEEE J Selected topics in applied earth observation and RS* 2 (1): 21-26.

White JC, Wulder MA, Varhola A, Vastaranta M, Coops NC, Cook BD, Pitt D, Woods M, 2013. A best practices guide for generating forest inventory attributes from airborne laser scanning data using an area-based approach. Natural Resources Canada, Canadian Forest Service, Canadian Wood Fibre Centre, Victoria, BC. Information Report FI-X-010, 39 pp.

White JC, Wulder MA, Hobart GW, Luther JE, Hermosilla T, Griffiths P, Coops NC, Hall RJ, Hostert P, Dyk A, Guindon L, 2014. Pixel-based image compositing for large-area dense time series applications and science. *Can J Remote Sens* 40 (3): 192-212. <https://doi.org/10.1080/07038992.2014.945827>

White JC, Coops NC, Wulder MA, Vastaranta M, Hilker T, Tompalski P, 2016. Remote sensing technologies for enhancing forest inventories: a review. *Can J Remote Sens* 42: 619-641. <https://doi.org/10.1080/07038992.2016.1207484>

White JC, Wulder MA, Hermosilla T, Coops NC, Hobart GW, 2017. A nationwide characterization of 25 years of forest disturbance and recovery for Canada using Landsat time series. *Remote Sens Environ* 194: 303-321. <https://doi.org/10.1016/j.rse.2017.03.035>

Wulder MA, 1998. Optical remote-sensing techniques for the assessment of forest inventory and biophysical parameters. *Progr Phys Geog* 22 (4): 449-476. <https://doi.org/10.1177/030913339802200402>

Wulder MA, Dymond CC, 2004. Remote sensing in survey of Mountain Pine impacts: review and recommendations. MPBI Report. Canadian Forest Service. Natural Resources Canada, Victoria, BC, Canada. 89 pp.

Wulder MA, Masek JG, Cohen WB, Loveland TR, Woodcock CE, 2012. Opening the archive: how free data has enabled the science and monitoring promise of Landsat. *Remote Sens Environ* 122: 2-10. <https://doi.org/10.1016/j.rse.2012.01.010>

Wulder MA, Hilker T, White JC, Coops NC, Masek JG, Pflugmacher D, Crevier Y, 2015. Virtual constellations for global terrestrial monitoring. *Remote Sens Environ* 170: 62-76. <https://doi.org/10.1016/j.rse.2015.09.001>

Wulder MA, White JC, Loveland TR, Woodcock CE, Belward AS, Cohen WB, Fosnight EA, Shaw J, Masek JG, Roy DP, 2016. The global Landsat archive: Status, consolidation, and direction. *Remote Sens Environ* 185: 271-283. <https://doi.org/10.1016/j.rse.2015.11.032>

Xie Q, Zhu J, Wang Ch, Fu H, López-Sánchez JM, Ballester-Berman JD, 2017. A modified dual-baseline PolInSAR method for forest height estimation. *Remote Sens-Basel* 9 (8): 819. <https://doi.org/10.3390/rs9080819>

Xie Y, Sha Z, Yu M, 2008. Remote sensing imagery in vegetation mapping: a review. *J Plant Ecol* 1 (1): 9-23. <https://doi.org/10.1093/jpe/rtm005>

Zald HSJ, Wulder MA, White JC, Hilker T, Hermosilla T, Hobart GW, Coops NC, 2016. Integrating Landsat pixel composites and change metrics with LiDAR plots to predictively map forest structure and aboveground biomass in Saskatchewan, Canada. *Remote Sens Environ* 176: 188-201. <https://doi.org/10.1016/j.rse.2016.01.015>

Zarco-Tejada PJ, Diaz-Varela R, Angileri V, Loudjani P, 2014. Tree height quantification using very high resolution imagery acquired from an unmanned aerial vehicle (UAV) and automatic 3D photo-reconstruction methods. *Eur J Agron* 55: 89-99. <https://doi.org/10.1016/j.eja.2014.01.004>

Zarco-Tejada PJ, Hornero A, Hernández-Clemente R, Beck PSA, 2018. Understanding the temporal dimension of the red-edge spectral region for forest decline detection using high-resolution hyperspectral and Sentinel-2A imagery. *ISPRS J Photogramm* 137: 134-148. <https://doi.org/10.1016/j.isprsjprs.2018.01.017>