

Table S2. Mean and standard deviation of the main attributes of the 14 shrubs species

Family	Species	n	DCH (cm)	HT (m)	CD (m)	CL (m)	NS (n)
Cortaderal site							
Elaeocarpaceae	<i>Aristotelia chilensis</i> (Mol.) Stuntz (Maqui)	21	3.60 ± 2.3	2.58 ± 0.9	1.81 ± 0.9	1.86 ± 0.9	4.48 ± 3.0
Salicaceae	<i>Azara dentata</i> Ruiz & Pavon (Corcolen)	42	4.85 ± 2.6	2.23 ± 0.9	1.49 ± 0.8	1.62 ± 0.8	4.40 ± 4.6
Asteraceae	<i>Baccharis macraei</i> Hook. & Arn. (Vautro)	21	2.34 ± 0.7	1.52 ± 0.4	1.45 ± 0.5	1.33 ± 0.4	4.67 ± 3.0
Berberidaceae	<i>Berberis horrida</i> Gay (Michay)	20	1.52 ± 0.6	1.54 ± 0.5	0.87 ± 0.3	1.28 ± 0.5	4.40 ± 2.1
Escalloniaceae	<i>Escallonia pulverulenta</i> (Ruiz et Pav.) Pers. (Corontillo)	23	3.46 ± 1.9	2.52 ± 0.9	1.23 ± 0.8	1.57 ± 0.8	2.70 ± 2.6
Asteraceae	<i>Gochnatia foliosa</i> D.Don (Mira Mira)	21	3.54 ± 1.8	1.99 ± 0.6	1.90 ± 0.8	1.60 ± 0.6	11.00 ± 7.9
Rosaceae	<i>Kageneckia oblonga</i> Ruiz et Pav. (Bollen)	21	3.93 ± 2.2	2.56 ± 1.1	1.08 ± 0.6	1.40 ± 0.7	1.19 ± 0.4
Proteaceae	<i>Lomatia hirsuta</i> (Lam.) Diels ex J.F.Macbr. (Radal)	20	4.88 ± 1.9	3.46 ± 1.1	1.64 ± 0.3	2.66 ± 0.8	2.65 ± 3.1
Myrtaceae	<i>Myrceugenia obtusa</i> (DC.) O.Berg (Arrayancillo)	23	6.03 ± 3.5	2.23 ± 0.8	1.79 ± 0.9	1.83 ± 0.7	5.91 ± 4.9
Lamiaceae	<i>Teucrium bicolor</i> Sm. (Oreganillo)	21	0.59 ± 0.3	0.90 ± 0.2	0.52 ± 0.2	0.67 ± 0.2	6.29 ± 3.2
Miraflores site							
Asteraceae	<i>Baccharis linearis</i> (Ruiz & Pav.) Pers. (Romerillo)	24	3.62 ± 1.8	1.48 ± 0.5	1.48 ± 0.7	1.14 ± 0.4	5.63 ± 4.1
Euphorbiaceae	<i>Colliguaja salicifolia</i> Gillies & Hook. (Colliguay)	23	3.64 ± 1.2	2.03 ± 0.7	2.82 ± 1.1	1.54 ± 0.5	13.00 ± 7.5
Asteraceae	<i>Podanthus mitique</i> Lindl. (Mitique)	22	2.30 ± 1.2	1.82 ± 0.5	1.71 ± 0.6	1.70 ± 0.5	10.82 ± 6.8
Rhamnaceae	<i>Retanilla trinervia</i> (Gillies & Hook.) Hook. & Arn. (Tevo)	20	6.00 ± 2.6	2.76 ± 0.8	3.37 ± 1.3	2.57 ± 0.8	7.35 ± 3.7

n = number of samples, DCH = diameter at collar height, HT = total height, CD = crown diameter, CL = crown length, NS = number of stems